

APPENDIX C

REVIEW OF HYDROGEOLOGY-RELATED INVESTIGATIONS IN THE UNITED STATES PART OF THE MESILLA BASIN REGION—1890 to 2010

The southeastern Basin and Range (B&R) physiographic province has been the center of major advances in the Earth sciences since the Civil War, with many pioneering contributions made by early “giants” of North American geology. The area of primary interest includes the Mexican Highland section of the B&R province and the southern Rio Grande rift tectonic province in New Mexico, Trans-Pecos Texas, and Arizona (e.g., Gile et al. 1981, Mace et al. 2001, Connell et al. 2005, Hawley 2005). It is located in “Ground Water Region 7—Central Alluvial Basins” as defined by Heath (1988), and described by Anderson and others (1988). Emphasis here is on the evolution of conceptual hydrogeologic-framework models of intermontane-basin and river-valley aquifer systems in the United States part of the Mesilla Basin region (**Fig. C1-1**) between 1890 and 2010.



Figure C1-1. Index map of the Mesilla Basin region (MBR) showing locations of the NW WRII Study Area (beige rectangle), major landscape features in the Mexican Highland section of the B&R province, and structural basins of the RG-rift tectonic province. Blue shading shows the approximate extent of the areas inundated by pluvial-Lakes Palomas and Otero at their respective Late Pleistocene high stands in the Los Muertos, and the Tularosa RG-rift basin areas. El Paso del Norte is shown as a yellow EPdN. Swanson Geoscience, LLC compilation on a 2017 Google Earth® image-base.

C1. PIONEERING-PHASE INVESTIGATIONS—1890 to 1945

The first part of this review comprises a series of illustrated short descriptions of seminal contributions of individuals and institutions with strong ties to the southeastern B&R province, especially studies that focused on geologic (geomorphic, hydrostratigraphic, lithofacies, and structural) controls on groundwater flow and hydrochemistry. Many subsequent advances in characterization of hydrogeologic systems in this arid/semiarid region have their conceptual roots in the cited pioneering work.

Following its establishment in 1879, the U.S. Geological Survey was the major institution contributing to hydrogeologic/geohydrologic investigations on a local to regional scale (e.g., Maxey 1979; Rabbitt 1989; Deming 2002, p. 8-11; Davis and Davis 2005; *cf.* Hawley and Kernodle 2008). Other early sources of information on the hydrogeology of the northwestern Rio Grande basin include reports by Hill (1891 and 1892b), Barker (1898), Follett (1898), Hill (1900), and Siebenthal (1910). Moreover, at least some alpine-glacial and periglacial features had now been identified in source highlands of the Rio Grande and Colorado River in the New Mexico-Arizona region (Stone, 1901).

Prior to Slichter's research on groundwater-flow in the Mesilla Valley-El Paso del Norte area, emphasis of work throughout the northern Rio Grande basin was on surface-water and stream-connected shallow aquifers (aka *subsurface waters*; e.g., Newell 1893, Follett 1898), and only one report dealt specifically with the Mesilla Valley (Barker 1898). Waring and Meinzer's (1947) "Bibliography and index of publications relating to ground water prepared by the Geological Survey and cooperating agencies (USGS Water-Supply Paper 992)," remains the most comprehensive document of its type on the history of groundwater-resource investigations in the United States between 1865 and 1945.

C1.1. Field-Based Research on the Rate of Movement of Underground Waters, by Charles S. Slichter (1864-1946)

Mathematician and geohydrologist Charles S. Slichter is internationally recognized for his pioneering laboratory and field investigations of "the motions of ground waters" (Slichter 1899, 1902, 1905a; Anderson 1995; Davis and Davis 2005). Moreover, in his "observations on ground waters of the Rio Grande valley (1905b)," Slichter made the first definitive quantitative analysis of inter-basin underflow discharge in the "Narrows of the Rio Grande near El Paso (**Figs C1-2a, b; cf. Fig. C1-7**)."

Slichter (1905b) introduced his work in the Paso del Norte (EPdN) area as follows:

[p. 9] An investigation of the underflow of the Rio Grande was begun in the latter part of August, 1904, at the narrows of the Rio Grande, a few miles above El Paso, Tex., where the stream flows through a narrow gorge of limestone. At this place is the site of the proposed Mexican-American international dam [**Fig. C1-2a**]. At the surface of the water the distance between the walls of the gorge is less than 400 feet. The dam site has been investigated by the International (Water) Boundary Commission, organized by the joint action of the American and Mexican Governments, and maps and reports concerning the proposed dam will be found in the Proceedings of the International (Water) Boundary Commission [IBC], vol. 2, page 277.

[p. 9] A brief reconnaissance at the site of the proposed international dam indicated that there could be no underflow of any magnitude at this point. The distance between the walls of the gorge is less than 400 feet, and the test borings made by the Mexican Commission in 1897 seemed to indicate that the maximum depth to bed rock is 86 feet. A cross section of the gorge, based upon the Mexican borings, is shown in fig. 2 [**Fig. C1-2b**]. In this diagram the vertical and horizontal scale are the same. A cross section of less than 40,000 square feet could not transmit a large volume of ground water even if other conditions were favorable. The highest velocity ever determined for ground water is about 100 feet in twenty-four hours, and assuming this maximum velocity at the above cross section, and a porosity of one-third in the water-bearing

sands and gravel, the daily discharge would be 1,333,000 cubic feet, or 15½ cubic feet per second. The gradient of the water plane at the narrows is but 3.8 feet to the mile, and all other indications point to a low rather than a high velocity. . . .




FIG. 1.—Topographic map of the gorge of the Rio Grande above El Paso Tex., where the underflow measurements were made.

Figure C1-2a. (Slichter 1905b, Fig. 1) First map of hydrogeologic-framework details in the “Narrows of the Rio Grande” near the northwestern end of Paso del Norte (*cf.* Part C1.2, Fig. C1-4; Darton 1933, p. 131). The north-trending dashed line shows the location of **Figure C1-2b** (Slichter 1905b, Fig. 2) and the sites of test boring for a (then) proposed “international dam (*cf.* APNDX H5.4 [Figs. H-2 and H5-3].”




Figure C1-2b. (Slichter 1905b, Fig. 1) Schematic cross-section depiction of shallow-subsurface hydrogeologic and hydrochemical relationships at the Paso del Norte location shown on **Figure C1-2a**. It represents the first cartographic product of this detail to be prepared in the RG-rift region, and Meinzer (1923b, Fig. 16) used it to illustrate the basic concepts of “underflow” and “underflow corridor.”

[p. 11-12] The material in the river bed at the site of the proposed international dam consists of sand and fine gravel with occasional layers of silt. No boulders were encountered in sinking the test wells and the borings were made with great ease. The ground waters in the sands of the gorge were found to contain a large amount of dissolved solids. At a depth of 10 feet the waters contained about 100 parts per 100,000 of common salt, and the quantity became larger as distance from the surface increased. Below a depth of 32 feet so much salt was present that it was very difficult to determine the rate of motion of the ground waters. At a depth of 42 feet the common salt in solution amounted to 1,340 parts per 100,000; at 60 feet it reached 1,700 parts per 100,000; in a 50-foot test hole it amounted to 1,720 parts per 100,000. The total solids dissolved in the ground water were proportionately as large as the amount of common salt, so that at a depth of 40 feet the water was about half as strong as sea water, and at a depth of 60 feet it was about 30 per cent stronger than ordinary sea water. In fig. 2 the positions of the principal test wells are shown. . . .

[13] The velocities of the ground water in the tests described above are undoubtedly the maximum velocities in the narrows of the gorge, since pains were taken to make the tests in the coarsest strata encountered in the borings. Layers of fine silt were frequently met with in putting down the test wells, which probably accounts for the stagnant condition of the water below the 35-foot level. These layers of silt are undoubtedly imbricated in such a way that movement of the deeper ground waters is impossible.

The total cross section in which the ground waters move is about 35 feet in depth and 325 feet in width and has an area of 11,200 square feet. If it is assumed that the porosity of the

material is one-third and the maximum velocity of the ground water is 3 feet per day, the total discharge through the gorge does not exceed 11,200 cubic feet per twenty-four hours, or about 0.132 cubic foot per second, or about 50 gallons a minute. This amount of underflow is entirely insignificant. It is obvious that on account of the enormous quantity of dissolved solids the underflow would be worthless, no matter what its magnitude might be.

C1.2. Development of the Bolson/Semibolson Conceptual Model by Cyrus F. Tolman (1873-1942)

Stanford University Professor of Geology and Mining, Cyrus Fisher Tolman, was a pioneer in the field of hydrogeology and was author of the first textbook on *Ground Water* (Deming 2002, p. 414-415; Tolman 1937). Based on long-term field observations in the American Southwest, he (1909) was the first to make a clear geomorphic and hydrogeologic distinction between depositional systems in aggrading intermontane basins with topographic closure (*bolsons*) and those that are open in terms of both surface and subsurface flow (*semibolsons*). He recognized three basic classes of lithofacies assemblages in the continuum of closed and open basin landforms: Piedmont-slope facies (e.g., coalescent alluvial-fan) are present along the margins of both basin types, while basin floors in topographically *closed* (*endorheic*) systems include alluvial flats that grade to terminal ephemeral-lacustrine plains (e.g., *playas*, *barreals* [*barrials*], and *salinas*). Floors of basins that are integrated with external (*open* or *exorheic*) surface-flow systems, in marked contrast, include alluvial flats and fluvial plains that grade to basin outlets (*cf.* Bryan 1938; **Fig. C1-8**).

The *closed* and *open*, intermontane-basin (*bolson*) landforms of the New Mexico region were major sites of definitive research on paleo-lake and ancestral-fluvial systems outside the Great Basin and Colorado River region (*cf.* Allen 2005, Connell et al. 2005, Hawley 2005). Of special importance were studies of Late Pleistocene lakes associated with USGS “Water-Supply” investigations in the Estancia, Wilcox (SE AZ), Tularosa, Animas-Cloverdale, Playas and Hachita basins (Meinzer, 1911, 1922; Meinzer and Kelton, 1912; Meinzer and Hare, 1915; Schwennesen, 1918). These pioneering studies of Late Cenozoic bolson and river-valley aquifer systems were closely linked to technological advances in deep-well drilling and the development of electric and (gas-oil) internal-combustion engines.

C1.3. Hydrogeologic Investigations in the Mesilla Basin Region by Willis T. Lee (1864-1926)

Willis Thomas Lee developed the earliest model of ancestral Rio Grande (ARG) evolution (1907b, **Fig. C1-3**), and was the first (1907a) to suggest that the Kilbourne and Hunts Hole maars (his Afton Craters) were the product of hydromagmatic-explosive processes. He (1907b) also played a key role in locating a dam at the Elephant Butte site for irrigation-water storage and flood control. Other seminal studies by Lee in New Mexico included work on evaporite-karst processes in the lower Pecos Valley (1925a), and he also led the first scientific surveys of the Carlsbad Cavern (Bat Cave) complex (1924, 1925b; Kues 2014a, p. 91-92).

The relatively advanced state of early conceptual models of Jornada (del Muerto) and Mesilla Basin hydrogeology is quite evident in the following excerpts from W.T. Lee’s (1907b) Water resources of the Rio Grande Valley in New Mexico: USGS Water-Supply Paper 188 (p. 10-11, 39-40). The accuracy of Lee’s interpretation of regional ancestral Rio Grande (ARG) evolution in the Mesilla Basin region is illustrated by **Figure C1-4** (Hawley et al. 2009, Fig. 9).



Figure C1-3. USGS geologist Willis T. Lee on his cutting-edge, AT&SF RR field vehicle during a Jornada del Muerto mapping expedition (*circa* 1904-1905; *cf.* Lee 1907b).

The following excerpts from “Water resources of the Rio Grande Valley in New Mexico: USGS, Water-Supply Paper 188” (p. 10-11, 39-40) illustrate the accuracy of Lee’s (1907b) interpretation of hydrogeologic system evolution in the southern Rio Grande-rift region:

CENTRAL AREA—PLAINS (Lee 1907b, p. 10-11)

The second plain, locally known as "La Mesa," lies in the southern part of the Rio Grande region west of Mesilla Valley, and extends from the vicinity of Las Cruces southward into Mexico. It is similar to the Jornada in many ways. Its altitude is the same as the southern end of the Jornada, and the two formed a single plain previous to the excavation of Mesilla Valley. La Mesa has a width of 20 miles or more and is undissected by erosion and entirely wanting in lines of surface drainage. It contains several broad shallow depressions, but, unlike those of the Jornada, these do not retain storm waters for any appreciable length of time. Although inclined slightly to the south] the surface appears practically level over an area of more than 1,000 square miles.

To a depth of at least 945 feet, the depth of the deepest well, the material in La Mesa consists of clay, sand, and rounded pebbles of quartzite, argillite, and a great variety of hard igneous and metamorphic rocks, with a subordinate amount of angular, debris. The surface is notably more sandy than that of the Jornada, and wells sunk in it encounter a greater proportion of fine material than occur in the Jornada.

In the northern part of La Mesa there are gravel beds of considerable size at the surface, but these become less numerous toward the south, until near the Mexican boundary sand alone is exposed, and the surface becomes practically level. The region was not explored south of the Mexican boundary for the purposes of this report, but from the summit of the Potrillo Mountains the sandy plain appeared to continue southward unbroken as far as the eye could reach. It is probable that La Mesa is the northern extremity of the broad interior basin of northern Mexico, the lowest parts of which, containing undrained lakes, occur 25 to 50 miles south of the international boundary. At some former time this basin was probably occupied by a large lake, the northern extremity of which covered La Mesa [Bolson de Los Muertos, and Lake Cabeza de Vaca of W.S. Strain 1966 and 1971; **Figs. C1-1 and C1-4**].




Figure C1-4. (Hawley et al. 2009, Fig. 9) Schematic depiction of Pliocene and Early Pleistocene evolution of the ancestral Rio Grande (ARG) fluvial-deltaic system in basins of the southern Rio Grande rift, with inferred age ranges. General area occupied by Upper Santa Fe Group ARG fluvial distributaries that spread out from a trunk-drainageway in the Palomas Basin above Hatch (dark pink), with a spill-out fluvial fan in eastern Mimbres Basin (light pink). ARG fluvial-plain remnants comprise the “La Mesa geomorphic surface.” The inferred Bolsón de Los Muertos ARG discharge zone is also shown (*cf.* Report **Parts 3.1.4** and **3.3.2**).

Lee also inferred that “the ancient Rio Grande flowed through the Jornada and La Mesa into the interior [Los Muertos] basin of Mexico, and that in comparatively recent geologic time changes occurred which turned it out of its valley [West Mesa area] and away from the interior [Los Muertos] basin toward the Gulf of Mexico (Lee 1907b, p. 22; **Fig. C1-2b**).” His observations on the hydrogeologic framework and groundwater-flow regime of the “LA MESA DISTRICT” are equally prescient.

UNDERGROUND WATERS—LA MESA DISTRICT (Lee 1907b, p. 39-40)

La Mesa district lies in the southern part of the Rio Grande region west of Mesilla Valley. Wells have been sunk in various parts of this district, both for railroad use and for stock purposes. No solid rock was encountered in any of the wells, most of which find water in abundance, but at a considerable depth, as indicated in Table 5 [**Tbl. C1-1**]. The deepest well on La Mesa, 945 feet, was bored by the Southern Pacific Railway Company at Lanark.* The company owns two other wells at the same place, one 648 feet and one 615 feet deep, the three yielding 50 gallons of water a minute. The material penetrated is sand with small waterworn pebbles, and contains water below a depth of 380 feet.

Table C1-1. (facsimile copy of USGS Water-Supply Paper 188, Table 5.—Records of bored wells in La Mesa district (Lee 1907b) Note that three wells (Herrington, Noria, and Potrillo) are on or near the former “El Paso and S.W. Rwy.” route, which is now the location of NM State Highway 9. The E.P. & S.W. and S.P. Rwy. Co. well are completed in ARG deposits (*cf.* Report **Part 3.4**).

Owner.	Location.	Total depth.	Depth to water.	Power used.	Material encountered.
		<i>Feet.</i>	<i>Feet.</i>		
Henry Brock.....	Sec. 30, T. 25 S., R. 2 W..	240	221	Gasoline..	Sand and waterworn gravels.
Do.....	Sec. 7, T. 24 S., R. 1 W...	430	386	do.....	Do.
Do.....	T. 24.....	515			Clay.
Mr. Hawkins.....	5 miles west of Picacho Mountain.	218	170	Gasoline..	
Robert Herrington.	2 miles northwest of Noria.	435	350	do.....	Sand and gravel.
J. F. Kilburn.....	T. 27 S., R. 1 W.....	^a 478	408		Do.
Do.....	6 miles northwest of Lanark.	388	370	Gasoline..	Sand and waterworn gravel.
S. P. Rwy. Co.....	Lanark.....	945	380	Steam.....	Do.
Lewis Bros.....	5 miles northeast of Lanark.	365	340	Gasoline..	Sand and gravel.
J. B. Stahling.....	10 miles west of Lanark.	^b 460	440		
Do.....	6 miles west of Lanark..	350	311	Gasoline..	
El Paso and S. W. Rwy.	Potrillo.....	240	220	do.....	Sand and clay.
Do.....	Noria.....	438	358	do.....	Sand and gravel.

^a 170 feet in bottom of crater.

^b 200 feet in bottom of crater.

Since the altitude of Lanark is 4,156 feet, the altitude of the water surface is 3,776 feet, while that at Bosque Seco, in Mesilla Valley, 15 miles northeast of Lanark, is 3,800 feet - 24 feet higher than at Lanark. At Noria, the altitude of which is 4,114 feet, the water surface, 358 feet below the surface of the land, is 3,756 feet above sea level. In the 12 miles between Lanark and Noria the water surface inclines to the south 20 feet, or at an average rate of 1.7 feet per mile. A line drawn through Bosque Seco, Lanark, and Noria would run somewhat west of the center of the old debris-filled valley of the Rio Grande for a distance of 27 miles. Along this line there is a fall of the water surface of 44 feet, or an average of 1.7 feet per mile. The gradient of the water table in Mesilla Valley between Bosque Seco (3,800 feet) and the southern end of Mesilla Valley (3,680 feet), a distance of about 32 miles, is 3.7 feet per mile. It is evident from these facts that the surface of the underground water has a regular gradient down the old channel through La Mesa, although it is less than the gradient of the river. A line of wells a few miles farther east in the center of the old valley would probably show a steeper gradient of the water plane.

The facts upon which the determination of gradient rests are not sufficiently numerous to make it conclusive. The depths to water determined and the indications that La Mesa is a part of the ancient debris-filled valley naturally leads to the inference that the course of the underflow should be southward through the detritus of La Mesa. It is possible, on the one hand, that additional data will show a gradient steeper than 1.7 feet per mile. On the other hand, it is possible that the original course of the underflow down the old channel has been reversed by reason of the down cutting of the river in Mesilla Valley and the accumulation of surface water in the gravels of La Mesa. The latter possibility is strengthened by the facts that La Mesa is nearly level and the material so porous that rain enters it without producing even temporary streams.

With respect to the future role of aerial photography in the earth sciences, and the constant need for ground truth, Lee also made these prescient observations (1922, p. 1-2):

The navigation of the air has accomplished much in many fields. Not only does it offer a new means of efficiency in [a variety of governmental], commercial and scientific pursuits. . .; but it has also opened a new world to the geographer, the photographer, and the geologist.

The ability of the camera to make instantaneous exposures and fix a clear image on a photographic plate enabled the [aerial] observer to obtain a record not only the scenes that he had viewed by also many that he might have missed . . . Immediately inventive genius was set to work to adjust the mechanisms of the camera to the demands of air photography and to prepare . . . films and sensitized paper necessary for the best results.

So satisfactory were the results and so great are the possibilities of further adaptation that there is an unfortunate tendency on the part of certain enthusiasts to make exaggerated claims that may retard progress. This is particularly true in the use of the air photograph in mapping. . . It cannot be reasonably expected to do away entirely the work of the ground surveyor. . . Observations from the air can never take the place of close examination of the ground, but it can be of great use in the location and study of land forms and geologic relations. Air photography is only an added means of obtaining information, although it promises to be a very important means.

C1.4. Early Advances in the Technology (and Art) of Drilling Deep Water Wells

Deep-well drilling in the 1890-1940 period was done very efficiently with cable-tool rigs such as the one at the Lubbock, TX National Ranch Heritage Museum shown in **Figure C1-5** (Bowman 1911; Meinzer and Hare 1915, p. 118).



Figure C1-5. Vintage cable-tool rig at the Lubbock, TX National Ranch Heritage Museum (*cf.* Bowman 1911, and Meinzer and Hare 1915, p. 118).

The following excerpt from Meinzer and Hare (1915, p. 118) illustrates the type of equipment that was used in drilling the deep wells that were required to supply the first steam railway operations throughout the binational Mesilla Basin region:

The valley [basin] fill is easily penetrated, and, except near the mountains where boulders are encountered, it presents few difficulties in sinking wells.

Most of the domestic wells are dug a short distance below the water table and are about 3½ feet in diameter. Some shallow wells have also been bored with augers propelled by hand. Where the water level is far below the surface, however, or where sinking to considerable distances below the water level has been necessary in order to obtain larger supplies of water that is less mineralized, machines propelled by horsepower, steam, or gasoline have been used.

The machine most commonly used for sinking these deeper wells is the portable standard [cable-tool] rig with percussion drill attached to a cable, the drill being withdrawn at intervals and the drillings removed by means of a bailer or sand bucket [Fig. C1-5; Bowman (1911) USGS WSP 257]. A machine of this type is among the most reliable for exploration work, especially where deep drilling is involved, and it is also well adapted for drilling through deposits containing boulders or hard layers. By its use water-bearing beds are generally detected, even though their yield is not great.

C1.5. New Mexico’s First “State Geologist:” N.H. Darton (1865-1948)

The field-based investigations of Nelson Horatio Darton (Kues 2014b) covered an enormous geographic and geologic-conceptual area (Darton 1899, 1905, 1914, 1916a-b, 1917, 1922, 1928a-b; Darton et al. 1916). His contributions to Late Cenozoic geology included formally naming the Ogallala Formation (1899, 1905). He was also a pioneer in the use of modern subsurface-mapping methods in his characterization of Luna County’s Mimbres basin aquifer system (1916a, Fig. C1-6).

Darton compiled the Deming folio for the Geologic Atlas of the United States series (1917), and other examples of these visionary synoptic coverages of geology, mineral resources, water supply, and soil conditions include the El Paso and Silver City Folios, respectively by Richardson (1909) and Paige and Darton (1916). Darton’s “Red Beds of New Mexico” (Darton 1928b) represents the first comprehensive map (a) and overview (b) of New Mexico geology. Of special importance in hydrogeological studies in the Mesilla Basin region (MBR), however, is his 1933 compilation of the first detailed-reconnaissance geologic map of the southern part of the Mesilla Basin and bordering mountain uplifts.

Like present-day NMGS Field Conference Guidebooks, the geologic railroad guides compiled by Darton and USGS associates (e.g., 1916, 1933) are invaluable sources of information not only on the local geologic and geomorphic setting, but also on its history and economy. Darton’s 1933 “Guidebook of the Western United States—Part F. The Southern Pacific Lines (USGS Bull. 845)” covers the present Union Pacific RR route that crosses the southern Mesilla Basin from the western Hueco Bolson and El Paso del Norte to the east-central Mimbres Basin (Fig. C1-7).




Figure C1-6. (Darton 1916b, Plate IX) First 3-D (fence-diagram) portrayal of lithofacies relationships in the shallow basin-fill aquifer system (Upper Gila Gp) in the Deming area of the central Mimbres Basin.




Figure C1-7. (facsimile copy of USGS Bulletin 845, SHEET 18) First reconnaissance geologic map of the Mesilla Basin region by N.H. Darton (1933). Bulletin 845 includes detailed commentary on geology and history along (1) the mainline of the Southern Pacific RR (now Union Pacific) between the El Paso, TX and Cambray, NM area, and (2) the southern (now abandoned) SPRR route between El Paso and Malpais Siding, NM (Darton 1933, p. 123-136, and p. 162-164; cf. Lee 1907b, Richardson 1909, Seager et al. 1987, Seager and Mack 1994, Seager 1995, Collins and Raney 2000, Hoffer 2001b).

Most descriptions of geology and hydrogeologic features in USGS Bulletin 845 retain their conceptual validity, as evidenced by the following extended passages, which start with entries on the El Paso area (elev. 3,725 ft/1,135 m; 1930 population about 131,500):

[p. 125] In 1852 a post office, called Franklin [after local land owner Franklin Coonze] was established here, and in 1859 the name was changed to El Paso. There was no town

development until after the Civil War—. . . . A few years after hostilities ceased a triweekly schedule was established for this region. The railroad reached El Paso from the east in 1883, and in the next few years brought many immigrants to the Rio Grande Valley. Since that time the city has had rapid growth.

El Paso has long been prominent as headquarters for the mining industry, although there are no notable mines in its immediate vicinity. For many years it has had the largest custom smelter* in the United States, . . . and treating [gold, silver, copper and other] ore[s] from New Mexico, Arizona, and Texas [and Mexico]. In 1930 the Nichols copper refinery was completed on the eastern edge of the city. Just west of this refinery are the Pasotex and Texas Co. oil refineries, which receive crude oil by a long pipeline from the Winkle field in Texas. Another pipeline brings gas to El Paso. The large cement plant on the western edge of the city furnished cement for the Elephant Butte Dam; it utilizes the limestone of the [Lower Cretaceous] Comanche series. Beaumont Hospital is a large Government establishment for tubercular soldiers, and Fort Bliss, 5 miles [8 km] northeast of the center of the city, . . . is the largest cavalry post in the United States. The Texas College of Mines [UTEP], a branch of the University of Texas, and Loretto College are also in El Paso. . . .

American Smelting and Refining Corporation (ASARCO) operation in El Paso del Norte (EPdN; USGS Bull. 845, PL. 18A). See discussion of environmental impacts in **APPENDIX H, Part 9.3.*

[p. 126] In New Mexico and Texas above El Paso the Rio Grande flows in a wide valley of alluvium [the Mesilla Valley], bordered by a high terrace plain [La Mesa surface]; At El Paso the valley is constricted to the narrow rock-walled pass that gives name to the city, but the bordering high terrace continues far down the valley. Below the pass the alluvial plain [El Paso/Juárez Valley floor] is a broad flat in which the river meanders widely, often changing its course by cutting new channels at time of freshets. The high terrace plain [western Hueco Bolson floor] that borders this valley terminates in bluffs and steep slopes, in places as much as 200 ft [61 m] above the bottom lands. The smooth [ARG-fluvial] plain at the top of these bluffs [La Mesa surface] extends far north as a wide bolson or desert flat between mountain ranges. Near El Paso there are several distinct benches, 3,800 to 3,950 feet above sea level, mostly in the form of mesas or projections from the base of the Franklin Mountains. These benches slope gently toward the river and are in part capped by caliche, an infiltration of calcium carbonate in the sand, which makes a material so hard that it helps preserve the tabular form and sharp edges of the mesas.

[p. 126-127] The Franklin Mountains form a high ridge on the southern prolongation of an axis of uplift which extends across central New Mexico from the Rocky Mountains. Probably this uplift is cut off to the south by a fault. The range rises abruptly about 3,000 feet [900 m] above the adjoining plains or valleys and culminates in Mount Franklin [north, alt. 7,192 ft/2,192 m]. The west side is mainly a dip slope of heavy beds of limestone with pronounced westerly dip; the east side shows many ridges, irregular lower crests, and buttes, and [is] deeply cut by canyons. The range is a typical tilted block of the basin-range type, which predominates in a large part of the Southwest.

[p. 128] The mountains to the south, in Mexico, are the Sierra Guadalupe [de Juárez] and the Cerro de Muleros [del Cristo Rey].

Wells in El Paso are to have penetrated valley fill [SFG basin fill] to a depth of 2,285 ft (696 m). Fossil bones found in Quaternary deposits have been determined as *Elephas Mammuthus] columbi*, *Equus complicatus*, and *Tapirus haysii?*, representing an elephant, an ancient horse, an a tapir, all of which have been extinct for many centuries [millennia]

[p. 128-129] The Southern Pacific [now Union Pacific] lines enter New Mexico on crossing the Rio Grande just west of El Paso, the State line being in midstream. . . .

[p.131] NORTH LINE FROM EL PASO, TEXAS TO MESCAL, ARIZONA

From El Paso westward to Tucson the Southern Pacific Railroad has two lines—one going by way of Deming and Benson [present UPRR] and the other (former El Paso & Southwestern Railroad [now abandoned]) by way of Columbus and Douglas [now route of NM Hwy. 9; **Fig. C1-7**; cf. Rpt. **PL. 9k**]. Leaving El Paso, the north line of the railroad follows the north [NE] bank of the Rio Grande for some distance, with Mexico in plain view on the opposite bank. In about 1 mile a large [ASARCO] smelter is passed, and in 2 miles a cement plant, near which are large quarries in limestones (of Comanche or Lower Cretaceous age) in a downfaulted block at the south end of the Franklin Mountains. It is the presence of this rock and a mass of intrusive [Campus Andesite] porphyry that cause the [EPdN] constriction of the river valley at El Paso (Spanish, the pass). At the entrance of this pass the railroad crosses the river into New Mexico and skirts the north side of Cerro de Muleros [del Cristo Rey], a high ridge which lies mostly in Mexico. . . .

[p. 131-132] The Cerro de Muleros consists of a mass of limestone, shale, and sandstone of [Lower] Cretaceous age penetrated and tilted by a large mass of [Eocene andesite] porphyry. The lower or quarry limestone in this succession is well exposed in the first railroad cut west of the river. It is overlain by nodular and slabby limestones and shales containing large numbers of Washita fossils and grading upward in to a thick mass of dark shale in which there are deep cuts extending to and beyond Brickyard siding. This shale is extensively worked for brick, hollow tile, etc., on the west bank of the river below the 2 railroad bridges. . . .

[p. 132] Just above Anapra siding [alt. 3,869 ft/1,179 m; pop. 27], where north [main SPRR] and south [now abandoned] lines are close together and joined by switches to be used in case of necessity, the railroad grade ascends the terrace of valley fill [Upper SFG], the edges of which margins the Rio Grande Valley in a long line of steep slopes. The top is attained near Strauss siding [alt. 4,110 ft/1,253 m]. Along the upgrade are many fine exposures of gravel and sand of which the terrace is composed. This material was deposited by the Rio Grande at an earlier stage of its history, when it flowed west of the Cerro de Muleros [Cristo Rey] and another high range to the south [Sierra de Juárez] and emptied into Guzman [Los Muertos] Basin in Chihuahua, Mexico. This was before its present course was developed through the “pass” at El Paso [cf. Lee 1907b, p. 10-11]

[p. 133] From Strauss siding the railroad goes northwest over the wide alluvial plain that extends entirely across the southwestern part of New Mexico. . . . The northerly trend of the railroad in this area is taken to avoid the large rugged area of volcanic rocks [basalt flows] of the West Potrillo Mountains and its extension to the north.

The thick body of sand and gravel underlying the plain has been drilled for water at several points along the railroad. A boring at Lanark [alt. 4,168 ft/1,270; pop. 40] passed through 950 ft (390 m) of beds, all supposed to be valley fill but possibly including some underlying Tertiary or Cretaceous strata [Lee 1907, Tbl. 5; Rpt. **TBL. 1**, no. 239]. It found water which rises approximately to the level of the Rio Grande Valley, 15 [8 mi/13 km] miles east. A boring at Kenzin [at eastern edge of Aden-Robledo Uplift], several [15 mi/24 km] miles beyond Lanark, passed through 550 feet [168 m] of clay and sand with water in its lower part and continued through [Lower Tertiary andesitic] rock 527 feet [161 m] farther.

Beyond Rutter siding great [basalt] lava fields are in sight to the southwest, and near Afton and Kenzin sidings [alt. 4,203 ft/1,281 m and 4,271 ft/1,302] the tracks skirt the edge of a fresh recent-looking [Aden/Afton] lava flow

[p. 134] Eight miles [13 km] southwest of Rutter siding are two large “holes” in the wide terrace plain [La Mesa surface] which have been the source of much wonderment for many years. They were originally called Los Corrales de Piedra (rock corrals). The more northerly, Kilbourne Hole, ..., is 300 ft [91 m] deep and is encircled with a rim of loose material 50 to 150 feet [15 to 46 m] high, so that in places it has a total depth of 450 feet [137 m],.... Hunts Hole, 2

miles [3.2 km] to the south, is closely similar but smaller and has a rim of less height.... The material in the wall of Kilbourne Hole (see fig. 28 [from Lee 1907a]) is stratified [Upper SFG-ARG] sand similar to that which underlies the wide surrounding [La Mesa] plain, capped by a 15-foot [5 m] layer of [basalt] lava, which thins out to the southeast. The encircling rim is composed mostly of soft sandstone, strongly cross-bedded and including some cinders, fragments of pumice, and many angular blocks of lava. A remarkable fact is that much of the cross-bedding slopes toward the hole. It is believed that these two holes were caused by a volcanic explosion probably with outburst of water. Steam doubtless accumulated in sand under the lava sheet until the pressure was sufficient to cause the explosion.

In the sand penetrated by a 170-foot [52 m] boring in the Kilbourne Hole, part of the jaw of a Pleistocene horse was found at a depth of 70 feet [21 m], possibly in material that caved from the sides of the hole. At the bottom of the boring a large accumulation of warm Sulphur water was found.

[p. 134] Through Kenzin, Pronto, Aden, Chappel, and Dona sidings lava field are in sight in every direction, especially on the south side of the railroad; most of them are recent outflows of scoriaceous basalt, but some are rhyolite [and basaltic andesite] of Tertiary age.

Aden Crater, 4 miles [6 km] south of Pronto siding and 7½ miles [12 km] southeast of Aden siding, is a cone of lava undoubtedly marking the vent from which came one of the large recent lava flows skirted by the railroad in this vicinity. In its top is a deep blowhole or steam vent in the lava, in which many animal skeletons have been found, including coyotes, bobcats, and other animals of the present fauna, and a remarkable ground sloth which Lull¹ has identified as *Nothrotherium shastense*. The remains were partly buried in bat guano in a sloping chamber about 100 feet [30] below the surface. The bones of the sloth were held together by the original ligaments, and some of the periosteum, patches of skin, muscle fibers, and claws remain. Most of the hide has been devoured by fellow victims, whose teeth marks are visible on the remaining fragments. The sloth and other animals had evidently fallen into this hole, which is a natural trap in the crater rim. The time was many thousands of years ago, for the sloth is a species found also in the Rancho La Brea asphalt in Los Angeles, where it occurs with bones of middle [and late] Pleistocene age.

¹Lull, R.S., 1929, *A remarkable ground sloth: Yale University Peabody Museum Memoir*, v. 3, part 2, 21 p.

Just south of Aden [siding, alt. 4,391 ft/1,338 m] is a prominent knob of rhyolite [Mount Aden] of the older volcanic series with a basalt flow at its foot, and 3 miles [5 km] west are other [p. 136] knobs of [Bell Top] rhyolite tuff capped by [Uvas] basalt of the older [Oligocene] succession. The volcanic area here [now in NE part of Mimbres River basin] extends so far north as to cause considerable deflection of the railroad to reach a long tangent that passes through Chappel [siding, alt. 4,279 ft/1,304 m] and extends [sub-parallel to I-10] beyond Carne [siding, elev. 4,193 ft/1,278 m].

At Chappel the lava fields are left behind, but detached igneous masses are in sight not far to the south and north. Two miles [3.2 km] south of Dona siding [near Doña Ana-Luna County Line] is the very prominent Providence Cone, rising nearly 300 feet [90 m] above the surrounding desert plain. . . . Far to the west from this place extends the level-floored Mimbres [River] Valley, which merges into the Lake Guzman [Bolsón de los Muertos] Desert, Mexico. This extensive basin has no surface outlet to the Rio Grande.

Near Cambray [alt. 4,229/1,289 m; pop. 40], the highway [now I-10] crossed the railroad; and after following the track for some distance, goes due west to Deming.

C1.6. Field-Based and Government-Administrative Contributions to Geohydrology and Hydrogeology by O.E. Meinzer (1876-1948)

Oscar Edward (O.E.) Meinzer is widely regarded as the “father of groundwater hydrology (Hackett, 1964).” His most significant contributions to hydrogeology were in two specialty areas: Basin and Range hydrologic systems, and federal-government service (Maxey 1979, Fryar 2007). Almost all of his early (1907-1912) fieldwork was done in the New Mexico region, but duties as Chief of the U.S. Geological Survey, Ground Water Division (1912 to 1946) occupied most of his professional life (Deming 2002, p. 86-87). It is important to note here that the first appointee to his 3-person staff was 24-year-old Kirk Bryan, a 1909 UNM geology graduate (*cf.* **Part C1.4**). Meinzer “connected” with the physical and cultural landscape of the New Mexico region at the very beginning of his career, so it seems reasonable to infer that many of his later contributions of national significance were strongly influenced by this early field experience. Major contributions to Basin and Range province hydrogeology are summarized in the following excerpt from Hawley (2014b, p. 117-118):

Meinzer’s first USGS publications dealt with the hydrogeology of closed intermontane basins of central New Mexico and southeastern Arizona. These “Geology and water resources” reports were released between 1911 and 1915, although most of the fieldwork was completed by 1912, and include Meinzer (1911) on the Estancia-Encino Valley area; Meinzer and Kelton (1912) on the Sulphur Springs Valley of southeastern Arizona (pluvial-Lake Cochise basin); and Meinzer and Hare (1915) on the Tularosa (pluvial-Lake Otero) Basin. They are still recognized as the seminal hydrogeologic studies on pluvial-lake basins in arid to semiarid parts of the American Southwest (Allen, 2005; Hawley, 2005; Hawley and Kernodle, 2008). Even cursory review of the many maps and photographs in these early papers shows how productive this pre-automotive era, Illinois “farm boy” could be when quality field work was facilitated by lots of equine and/or *shanks-mare* assistance.

Following additional field studies in the Big Smoky Valley area of east-central Nevada, Meinzer synthesized his “pluvial-lake” research in a 1922 paper *Map of the Pleistocene lakes of the Basin and Range province and its significance*. This is the first lucid description of “the extent to which ancient lakes would be restored by lowered temperatures (Waring and Meinzer, 1947, p. 193).” He also proposed that hydrologic conditions conducive to lake formation in the cool-dry northern Great Basin region were valid modern analogs of Late Pleistocene paleohydrologic systems in the relatively warm-dry playa-lake basins of the southeastern Basin and Range province. In addition, the Meinzer and Kelton (1912) Sulphur Springs Valley report exemplifies a pioneering “integrated, joint investigation of chemical and physical hydrogeology (Meyer et al., 1988, p. 5). While the above-cited publications relate to Meinzer’s direct contributions to geologic research in the New Mexico region, they also help explain his future effectiveness in federal-governmental science and administration, particularly when prior Midwestern rural-life and educational experiences are taken into account.

C1.7. Hydrogeologic Contributions by Albuquerque Native, Kirk Bryan (1888-1950)

Albuquerque native and 1909 UNM graduate, Kirk Bryan (1888-1950) is now primarily recognized for his contributions to geomorphology and geoarchaeology (Sharp 1993, Leopold 2004). From all perspectives, Bryan, his Harvard students (including Luna B. Leopold), and many professional colleagues have been major figures in the fields of geomorphology and Late Cenozoic geology since the early 1920’s. He was a major contributor to the development of current models of hillslope-landform evolution, and many other aspects of physiography in both semiarid and humid regions (Bryan 1923, 1925, 1936, 1940a-b, 1941; Bryan and McCann 1937, 1938). Much of his career, however, was devoted

to field investigations in Arizona and New Mexico (Hawley 2005; Hawley and Kernodle 2000, 2008; Hawley et al. 2001; Phillips 2014).

Bryan joined O.E. Meinzer's new USGS Division of Ground Water in 1912, and his important hydrogeologic contributions include development of the first synoptic conceptual model of basin-fill aquifer systems in the Rio Grande rift tectonic province. This work is summarized in the "Geology and ground-water conditions of the Rio Grande depression in Colorado and New Mexico" (Bryan 1938) for a "[US] National Resources Committee . . . Regional Planning" publication, which also included C.V. Theis' first detailed review of "ground water in the middle Rio Grande valley" (*cf.* Theis 1935a, b; 1938).

As earlier done by W.T. Lee (1907b), Bryan recognized that the river-linked series of deep structural basins, which extend from southern Colorado to Trans-Pecos Texas, form a unified geologic and geohydrologic system. This regional tectonic feature, which Bryan initially named "the Rio Grande depression," is now designated the Rio Grande rift (RG-rift; Kelley, 1952; Chapin, 1971; Chapin and Seager, 1975; Hawley 1978; Riecker 1979; Keller and Cather 1994). Bryan's (1938) major contribution to upper Rio Grande basin evolution was his observation that (p. 205): "The main body of sedimentary deposits of the Rio Grande depression, from the north end of the San Luis valley to and beyond El Paso, is considered to be the same general age and to belong to the Santa Fe formation...." In general, the basins appear to have been elongated into ovals and to be divisible into two major types: basins with a through-flowing river and basins with enclosed drainage." He also noted (p. 221) that basins of the Rio Grande depression "differ from other basins [in the B&R province] principally in being strung like beads on a string along the line of the Rio Grande."

In comparison to Tolman (1909 and 1937), Bryan's 1938 conceptual model of intermontane-basin hydrogeologic systems was developed in much greater detail and presented in a semi-quantitative format. **Figure C1-8**, adapted by Hawley and Kernodle (2008) from Bryan (1938, Figures 51 and 52), clearly demonstrates a basic understanding of integrated surface and subsurface flow regimes in "completely enclosed" and "incompletely enclosed" basins of the "Rio Grande depression," and similar geologic terranes throughout the Basin and Range province. This robust model of groundwater flow systems in a variety of *bolson* and *semibolson* hydrogeologic settings was originally used in implementing the 1938 Rio Grande Compact, and in continues to serve as a definitive, basin-scale geohydrologic model.




Figure C1-8. Conceptual model of geologic and hydrologic relations in intermontane basins of the Rio Grande “depression.” *From* reformatted version of Bryan (1938) *in* Hawley and Kernodle 2008, Figs. 51 and 52).

C1.8. Exploration Geophysics and Hydrogeology in the El Paso Area

Notable USGS contributions to early groundwater-resources studies in the El Paso area include the pioneering work of A.N. Sayre and E.L. Stephenson (1937) on the use earth-resistivity geophysical-survey methods to locate “salt-water bodies” in local basin-fill aquifers. Another seminal “ground-water” investigation by Sayre and Penn Livingston, which was completed before World War II, was done in cooperation with the El Paso Water Board and the Texas State Board of Water Engineers. Their 1945 Water-Supply Paper (919) includes descriptions of geology, ground-water occurrence, and available water-quality information. B.J. Hibbs (2/15/2021 written communication) states that (1) the “first transboundary water table map was also generated by Sayre and Livingston (1945, Fig. 4),” and (2) it was based on data “compiled in Sayre and Livingston (1945, appendices labeled Well Logs, Records of Water Levels in Observation Wells in El Paso Area, Deep Wells Analyses of Water, and Records of Wells in the El Paso Area.” He notes that: “While hydrogeologic problems from aquifer overdraft took more time to develop than predicted by Sayre and Livingston (1945), their report set in motion accelerated inventories of groundwater resources in the United States and Mexico and established beneficial data exchanges.”

C2. MULTI-INSTITUTION HYDROGEOLOGIC/GEOHYDROLOGIC INVESTIGATIONS—1945 to 1971

Throughout the United States, the WW II and early post-war period witnessed the development of strong collaborative ties between geologists and hydrologists involved in groundwater investigations. In the New Mexico–Trans-Pecos Texas region, this was primarily due to the continued leadership role played by the USGS-Water Resources Division (WRD), NM Office of State Engineer, NM Bureau of Mines & Mineral Resources at NM Tech, Texas Water Commission, and (after 1964) the NM Water Resources Research Institute at NMSU (e.g., Conover 1954, Knowles and Kennedy 1958, Reeder 1957, Leggat et al. 1962, Spiegel 1962, and Spiegel and Baldwin 1963, Trauger and Doty 1965, Cliett 1969, Hawley 1969a, King et al. 1971, Trauger 1972). Report **Figure 1-4** is an index map of the Mesilla Basin Study Area that shows locations of groundwater-basins, hydrogeologic cross-sections (A-A’ to S-S’), population centers, and major landscape features, including the Mesilla Valley of the Rio Grande.

C2.1. USGS Studies in the Rincon and Mesilla Valley Areas—1946 to 1954

Figure C2-1 illustrates the initial developmental stage of potentiometric-surface maps that schematically depict GW-flow trends in the United States part of the Mesilla Basin region. This 1954 map by Clyde Conover (Plate 1) shows approximate 1947 water-level contours in central and southern Doña Ana County, New Mexico. Estimated depths to water in the 300 to 400-ft and > 400 ft ranges are shown, respectively, with light greenish yellow and green dotted patterns. This and subsequent maps comprise an invaluable history of ground-water flow evolution during the past 75 years (*cf.* King et al. 1971, Wilson et al. 1981, Frenzel and Kaehler 1992, Nickerson and Myers 1993, and Teeple 2017a).

C2.2. Cooperative Investigations in Lower Mesilla Valley Area—1952 to 1962

The USGS, Texas Water Commission, and City of El Paso initiated a cooperative study in 1952 that was designed to determine the quantity and quality of groundwater available in the Lower Mesilla Valley for (1) public supply of El Paso, and (2) industrial and irrigation use (Leggat et al. 1962). The study by Edward Leggat and associates, which was completed in 1958, included a number of innovative hydrogeological, geophysical, and hydrochemical practices with respect to groundwater-resource investigations in the Mesilla Basin region:

1. Application of standard “oilfield” borehole-geophysical logging methods in ground.
2. Drill-stem hydrochemical sampling of representative hydrostratigraphic intervals in most wells.
3. Formal introduction of the Santa Fe Group (SFG) lithostratigraphic concepts to basin-fill stratigraphy.

4. Subdivision of the SFG into informal “upper, medium [middle], and lower” hydrostratigraphic categories based on drill-cutting and geophysical-log analyses from deep test- and production wells (Figs. C2 and C2-3a, b).




Figure C2-1. (Report **Plate 9A extract**) Facsimile copy of central and southern parts of Plate 1 in Conover (1954). The map shows approximate 1947 water-level contours in central and southern Doña Ana County, New Mexico. Estimated depths to water in the 300 to 400-ft and > 400 ft ranges are shown, respectively, with light greenish yellow and green dotted patterns.




Figure C2-3. (modified from Hawley and Kernodle, 2000, Fig. 6) Correlation diagram of major Time-Rock (chronostratigraphic) classes, lithostratigraphic, allostratigraphic, and hydrostratigraphic units of Cenozoic Age in the southern RG-rift region (*cf.* NACOSN 2005; **APNDX. G**). Bedrock units: Qb–Quaternary basalt, Tb–Tertiary mafic volcanics, and Tv–older Tertiary intermediate and silicic volcanics, and associated plutonic-igneous and sedimentary rocks.




Figure C2-3b. (facsimile copy of Leggat et al. 1962, Fig. 5) “Geologic section B-B’, lower Mesilla Valley” that is located near the eastern end of Hydrogeologic Section I-I’ (32° N latitude alignment). Information on listed wells is on the **TABLE 1** spreadsheet: Q63-NA, Q172-no. 336, Q189-no. 335, Q72-no. 340, and Q73-no. 344 (*cf.* **Figs. C2-4** and **C2-5b**).

The following passages from the Leggat and others report (1962, p. 10-16) document the scope of this seminal contribution to the hydrogeology of the Mesilla Basin region. Detailed information on specific wells is on the Report **TABLE 1** spreadsheet (e.g., Q178-no. 329):

Santa Fe Group

The Santa Fe group underlies the lower Mesilla Valley and is exposed in nearly all the arroyos between the flood plain and the Franklin Mountains and in the bluffs at the east edge of La Mesa. In the uplands east of the Rio Grande, the top of the Santa Fe, mostly coarse sand and gravel containing some caliche-cemented boulders, probably is correlative with the Pleistocene cap of the Santa Fe, as defined by Spiegel and Baldwin (in preparation [1963]). A thick series of reddish to brown silty clay, fine to medium sand or poorly consolidated sandstone, and thick-bedded conglomerate underlies the coarse sediments.

Characteristic responses in electric logs of eight wells northwest of Canutillo suggest that the Santa Fe may be subdivided into two units (See Figures 4 and 5 [**Figs. C3-2a** and **C2-3b**]). In electric logs, the curve on the right side of the base line represents the relative resistivity of the individual beds. A deflection of the resistivity curve to the right (increase in resistance) usually is indicative of a fresh-water-bearing sand. Sand beds containing brackish or salty water have low resistance and cause little or no deflection of the resistivity curve. It is not possible to determine on the basis of available data if the units can be correlated over a large part of the valley.

The lower unit (the deep aquifer of the well field northwest of Canutillo) is composed of unconsolidated fine to medium sand; the percentage of clay is smaller than in the upper unit. The low, but uniform, resistivity response in the well field suggests that the sand in the lower unit is uniform, thick bedded, and relatively free of interbedded shale or clay [HSU-LSF, LFA 4—Rpt. **Fig. 4-3, Tbls. 4-1 and 4-2**]. The lower unit reaches a maximum thickness of at least 1,000 feet in well Q-178 [**no. 328**]; electric logs indicate that the lower unit thickens to the north and west. The spontaneous-potential curve (on the left side of the base line) of well Q-144 [**no. 281**] suggests that the lower unit in the south end of the valley is much thinner.

The upper unit of the Santa Fe, which contains the medium [Middle SFG] aquifer and a part of the shallow aquifer of the El Paso well field, is exposed in arroyos above the flood plain and in the bluffs of La Mesa. It consists of alternating layers of varied thickness of fine to coarse sand, gravel, and reddish-brown silty clay. Locally the sand is crossbedded, lenticular, and predominantly medium-grained. The clay is evenly bedded in many exposures along Mesa Road (U. S. 80), indicating that clay layers may extend laterally for considerable distances. In [Beneath] the flood plain, the upper unit reaches a depth of 470 feet in well Q-86 [NA], 678 feet in well Q-178 [**no. 328**], and possibly 1,100 feet in well Q-144 [**no. 281**]. According to electric logs and drillers' logs, the maximum thickness of the upper unit is at least 1,000 feet and perhaps as much as 1,400 feet.

Electric logs and drillers' logs reveal that in [beneath] the flood plain a thick-bedded limestone conglomerate underlies the lower unit but that east of the flood plain the conglomerate occurs within the Santa Fe. The conglomerate, derived from limestone that probably is of Cretaceous age (Virgil Barnes oral communication, Feb. 27, 1957), has a maximum observed thickness of 160 feet in well Q-173 [no. 327]. According to the electric and driller's logs of well Q-173 the conglomerate is underlain by clay, sand, and gravel. Although the sediments underlying the conglomerate probably are Santa Fe in age [here interpreted as pre-Santa Fe], they are not included in the lower unit. Figures 4 and 5 show that the conglomerate lies at a greater depth north and west of well Q-173 [**Figs. C2-5a and 5b**]. Well Q-178 [**no. 328**], depth 1,705 feet, did not penetrate the conglomerate. The highly resistive zone between 1,280 and 1,380 feet in well Q-178 is an igneous sill underlain by sand, shale, and clay of the lower unit of the Santa Fe. The igneous rock penetrated by well Q-178 may be associated with the series of volcanic rocks interbedded with the Santa Fe group of sediments near Las Cruces, New Mexico [interpreted as pre-Santa Fe bedrock-unit TIs herein].

The maximum thickness of the Santa Fe group in the lower Mesilla Valley is not known; unconsolidated sediments were penetrated to a depth of 1,705 feet in well Q-178 [**no. 328**]. The original thickness of the unconsolidated sediments in the area, the surface being about 300 feet lower than La Mesa, may have been about 2,000 feet. However, as the sediments thicken toward the center of the basin, it is probable that the maximum thickness of the Santa Fe is considerably greater.

The medium and deep aquifers of the Santa Fe group [SFG] are the major sources of ground water for public supply in the valley. Many wells obtain water from both the Santa Fe and the alluvium. Yields as large as 3,000 gpm of fresh water have been obtained from the Santa Fe north and northwest of Canutillo. South and southwest of Canutillo, the water in the Santa Fe is brackish or salty. Small to moderate quantities of fresh water have been obtained from the Santa Fe in the upland east of the Rio Grande, where the saturated thickness is much less.

C2.3. Digital Conversion of Borehole Geophysical and Drill-Cutting Logs—1987 to 1992

The seminal work of Leggat and others (1962), as illustrated in **Figures C2-2a and C2-2b**, continues to provide a robust model for the hydrostratigraphic and lithofacies interpretations of later workers. As noted in **Part C5.2**, hydrogeologic investigations between 1983 and 1997 included

collaborative efforts that involved the NMBMMR Environmental Geology and NM Tech Hydrology programs, the USGS-WRD (NM), the El Paso Water Utility (EPWU), and the NMOSE. Emphasis was on: (1) manual conversion of analog data from original borehole-electric logs into digital format, (2) compilation of drill-stem-sample information on water quality, (3) detailed petrographic analyses of drill cutting, and (4) interpretation of this subsurface database as an initial step in hydrogeologic-framework characterization in the Mesilla GW basin (e.g., Hawley and Lozinsky 1992, Nickerson and Myers 1993). The examples of this work from Hawley and J. Kennedy (2004) that are presented in **Figures C2-3 to C2-5**, illustrate the value of continuing to utilize the historic databases developed by Leggat and other (1962) and their contemporaries in the Mesilla Basin/Hueco Bolson region (*cf.* Knowles and R. Kennedy 1958).




Figure C2-4. (Report **Fig. 2-3**) NM WRRRI Study Area index map on a **PLATE 1** Hydrogeologic Map base. The Mesilla, Southern Jornada, and El Parabién groundwater (GW) basins (MeB, SJB and EPB) are outlined in green, orange, and red, respectively. Also shown are locations of major terrain features (including the Mesilla Valley, Selden Canyon and El Paso del Norte of the Rio Grande), hydrogeologic cross-sections **A-A'** to **S-S'**, and the El Paso/Ciudad Juárez and Las Cruces metropolitan districts.




Figure C2-5a. (Hawley and Kennedy 2004-Appendix A, Plate A7) Down-valley hydrogeologic cross-section of the lower Mesilla Valley that follows the approximate trends of (1) the Section O-O'-O''-O''' series (PL 50), and (2) Geologic section A-A' in Leggat and others (1962, Fig. 4; cf. **Fig C2-3a** caption). Detailed information on specific wells is on the Report **TABLE 1** spreadsheet: Vado test-no. 200, Berino test-no. 227, Colquitt Farms-no. 332, Q178 Howell-no. 328, Q197-no. 329, Q63-NA, Q202-no. 339, Q144-no. 279, and Lizard 2-no. 291. Drill-stem water-sample locations are shown with black rectangles, and the upper and lower numbers show, respectively, depth to top of screen in feet and sample TDS in mg/L. Borehole resistivity-logs (red: short-normal, blue: long-normal).




Figure C2-5b. (eastern part of Plate A3 in Hawley and Kennedy 2004-Appendix A) Transverse hydrogeologic cross-section of the lower Mesilla Valley that is located near the eastern end of Hydrogeologic Section I-I' (near 32° N alignment; cf. Fig. C2-4). See Fig. C2-3a caption, and Report TABLE 1 entries for additional information on specified test wells: La Union (MT-1)-no. 242, Q43-no. 322, Q178-no. 328, Mountain Pass-no. 333, La Tuna (Fed. Prison)-no. 332, and Q25-no. 330. Drill-stem water-sample locations are shown with black rectangles, and the upper and lower numbers show, respectively, depth to top of screen in feet and sample TDS in mg/L.

C2.4. Cooperative Investigations with El Paso Water Utility—1963 to 1970

The above-described Mesilla Valley investigations were primarily conducted by USGS hydro-scientists, and they temporally overlapped with studies in the Hueco Bolson of similar scope that were

covered in Water Supply Paper 1426 (Knowles and Kennedy 1958). For the next four decades (through the 1990s) Thomas E. (Tom) Cliett of the El Paso Water Utility (EPWU) played a major leadership role of geology-based groundwater investigations in the Texas part of the southern Mesilla Basin and western Hueco Bolson part of the Paso del Norte region (e.g., Mattick 1967, Cliett 1969, Knorr and Cliett 1985, Cliett and Hawley 1996; cf. Hawley et al. 2009). Cliett (1931-2015) graduated from Texas Western College in 1958 with B.S. in geology; and he immediately began a 32-year tenure with the EPWU groundwater-geology program. It soon expanded under his leadership with innovative aquifer exploration and development methods that included borehole geophysics, hydrochemical sampling, and ultimately aquifer-storage and recovery. The following quote from Cliett (1969, p. 210-212), illustrates his early grasp of the essential hydrogeological and hydrochemical components of the GW-flow system in the lower Mesilla Valley between Anthony, NM/TX and El Paso del Norte (**Fig. C2-6**):

Production in the Canutillo Well Field, which supplies the City of El Paso with a large portion of its water, is derived from three zones known as the shallow, intermediate, and deep which probably constitute only a two-aquifer system. Most of the shallow zone is made up of coarse sand, gravel, silt and clay and probably does not exceed 90 feet in thickness when referring to Recent alluvium, but the production zone itself extends to an average depth of about 150 feet. There is often no distinct lithologic break (as indicated by well cuttings) between the Recent [Late Quaternary] alluvium and the underlying Santa Fe Group basin fill within the shallow aquifer zone, and the two rock-stratigraphic units are hydraulically connected. The separation of the shallow and intermediate zones is based primarily on groundwater quality changes even though static levels in the intermediate average about 40 feet from the ground surface and only 10 feet in the shallow. There are no known continuous barriers between the two, but overlapping clay lenses possibly create a partial barrier to vertical components of groundwater movements.

The differing qualities of groundwater in the various zones may be noted in Figures 6 and 7 [e.g., **Figs. C2-7** and **C2-8**] and are characteristic of the Canutillo Well Field. However, the quality of the groundwater in the shallow zone becomes much better to the north and west of the field.

The deep aquifer is a fine-grained homogenous body of unconsolidated sand with superior quality groundwater and an artesian head of about 440 feet (190 psi). The aquifer is encountered at an average depth of about 500 feet which places the existing piezometric surface at 60 feet below the surface of the ground. Its average thickness is about 500 feet and its only known occurrence is in the Canutillo Well Field. An equally extraordinary anomaly, besides its excellent quality, is the temperature of the water of 98° F. [36.7°C].

The aquifer may represent a fluvial environment of an isolated nature where the ancestral Rio Grande acquired stream velocity magnitudes great enough to carry silts and clays to other points of deposition, leaving the fine-grained sand in its present locale.

King and others (1969) have speculated that the aquifer may represent deposition in a deltaic environment. It is also possible that the unit is basically an aeolian deposit analogous to the Medanos de Samalayuca dune complex in north-central Chihuahua (John Hawley, personal communication, July, 1969). In any case, the aquifer has rapid communication with a recharge zone of a highly permeable nature which is indicated by its water quality, and the recharge is directed through a zone with a high geothermal gradient.

Much more subsurface data are needed to make more accurate assumptions but undoubtedly future test drilling will prove a more extensive occurrence of the deep aquifer zone.




Figure C2-6. (facsimile copy of Cliett 1969, Fig. 5; reproduced with New Mexico Geological Society, Inc. permission) Index map of the lower Mesilla Valley showing locations of local geomorphic and cultural features, and hydrogeologic cross-sections C-C' and D-D' (Figs. C2-7 and C2-8, respectively). "RIMROCK" delineates the Mesilla Valley-West Mesa (La Mesa surface) boundary and the constructional top of Upper Santa Fe Group fluvial-facies (ARG) deposits (USF-HSU2/LFA2&1).




FIGURE 6.
Generalized geology and groundwater occurrence in Canutillo Well Field.

Figure C2-7. (facsimile copy of Cliett 1969, Fig. 6; reproduced with New Mexico Geological Society, Inc. permission) Transverse “Generalized” geologic section C-C’, lower Mesilla Valley” that is located near the eastern segment of cross-section J-J’ (Fig. C2-6, PL. 5j).




FIGURE 7.
Generalized geology and groundwater occurrence, lower Mesilla Valley to El Paso Canyon.

Figure C2-8. (facsimile copy of Cliett 1969, Fig. 7; reproduced with New Mexico Geological Society, Inc. permission). Longitudinal hydrogeologic section D-D' of the lower Mesilla Valley between Anthony, TX and Sunland Park, NM. Cross-section location (**Fig. C2-6**) also approximates that of section A-A' in Leggat and others (1962, Fig. 4; **Fig. C2-3a**) and Plate A7 in Hawley and Kennedy 2004-Appendix A (**Fig. C2-5a**). Sedimentary units that underlie the “DEEP AQUIFER and intertonguing conglomeratic deposits of the Lower SFG hydrostratigraphic unit (HSU-LSF) are here correlated with the pre-Santa Fe (T1s) bedrock sequence of Early to Middle(?) Tertiary Age (Report **TABLE 2**). “INTERMEDIATE ZONE” deposits are primarily part of the Middle Santa Fe HSU, while the SHALLOW ZONE comprises Upper SFG basin fill with a thin (<100-ft, 30-m) cover of river-valley alluvial fill beneath the Rio Grande floodplain and bordering valley sideslopes.

For more than a decade following his retirement, Tom Cliett was still actively involved as a private consultant in groundwater exploration and well/wellfield design. Throughout his career he was an enthusiastic collaborator in all phases of aquifer-system characterization throughout the binational Hueco Bolson-Mesilla region, especially in the planning and implementation of USGS-NM OSE-NM Bureau of Mines collaborative studies completed in the early 1990s (**Part C2.3**). The following excerpt and diagrams from Cliett (1969, p. 210-214) illustrates some of the contributions of a consummate public servant, who devoted a lifetime to getting a sound grasp on the applied hydrogeology of the Paso del Norte region (**Figs. C2-5 to C2-7**). A cross-section diagram from Cliett and Hawley (1996, Fig. 5; **Fig. C2-9**) is also included to better illustrate Tom Cliett's significant contributions in the areas of groundwater chemistry, and hydrogeologic-framework characterization:

The geology of the Mesilla bolson is similar to that of the Hueco bolson and the basin fills are contemporaneous. By early Pleistocene (Kansan time) basin filling in the El Paso and Lower Mesilla Valley areas reached its climax (Kottlowski, 1958a; Hawley, et al. 1969; King, et al. 1969). The ancestral Rio Grande entered the Hueco bolson through Fillmore Gap [Pass] between the Organ and Franklin Mountains and was diverted to its present course in late mid-Pleistocene time (Strain, 1966).

Groundwater produced for irrigation and municipal purposes in the Lower Mesilla Valley occurs principally in deposits of the Santa Fe Group with many shallow wells completed in Recent [Late Quaternary] river alluvium.




Figure C2-9. (facsimile copy of Cliett and Hawley 1996, Fig. 5) Transverse hydro-chemical/stratigraphic cross-section D-D' is located near the eastern end of Hydrogeologic Section I-I' (32° N alignment, **Fig. C2-3**). "DEEP and INTERMEDIATE" SFG aquifer zones are, respectively, approximate equivalents of the Lower and Middle Santa Fe Hydrostratigraphic Units (LSF and MSF—**Fig. C2-2b**). Numbers to right of well symbols are water-quality values in mg/L tds. Detailed information on the following wells is on the Report **TABLE 1** spreadsheet: Anthony Test 3-no. **324**, UTV 5 (Q25)-no. **330**, La Tuna (Fed. Prison)-no. **332**, and Anthony Test 2-no. **351**. Note that the Anthony Test Wells are near Q178 (Howell)-no. **328** (**Figs. 2-8a**)

C3. USDA-SCS SOIL SURVEY INVESTIGATIONS DIVISION, NMSU (NM WRRI AND EARTH SCIENCE DEPT.) and NM TECH (NM BUREAU OF MINES & MINERAL RESOURCE DIV.) COLLABORATIONS —1964 to 1977

C3.1. Research Related to the USDA-SCS Desert Soil-Geomorphology Project


From October 1962 to May 1977, Hawley's geology-related activities in the New Mexico-West Texas region were performed as an employee of the Soil Survey Investigations Division of the USDA Soil Conservation Service, with administrative headquarters in Washington D.C. (Hawley 2005, p. 15-16). Until June 1971, most work was done from a field headquarters based at NMSU-Las Cruces, and during the following four years, field research was based at Texas Tech-Lubbock (1971-1974), and NM Tech-Socorro (1974-1975). Soil-geomorphic and hydrogeologic studies in Chihuahua were completed in 1968 and 1969 in cooperation with Mexican federal agencies (**APPENDIX D, Parts D2 and D3; and APPENDIX H**). The scope of geology-related work in the binational Mesilla Basin region (**Figs. C1-1**) is described in the following excerpt from the 26th NMGS Field Conference "Guidebook to the Las Cruces Country" (Hawley 1975b, p. 183):

The Desert Soil-Geomorphology Project refers to a Soil Conservation Service [USDA-SCS] investigation, between 1957 and 1972, of landscape evolution and soil development in a 400 sq. mi. area of Dona Ana County, New Mexico (Fig. 1 [**Fig. C-14**]). Its primary purpose was to gather basic information on soil-geomorphic relationships that would lead to increased accuracy and efficiency of the Soil Survey program in arid and semi-arid regions of the western states. A team of soil scientists and geologists from the S.C.S. Soil Survey Investigations Division . . . staffed the project. Research geologist R. V. Ruhe conducted initial geologic-geomorphic field studies from 1957 to 1960 and headed the project from 1957 to 1965 [*cf.* Ruhe 1962, 1964, 1967]. F. F. Peterson and J.W. Hawley were responsible for geomorphic and geologic research from 1960 to 1962 and 1962 to 1972, respectively. L. H. Gile was in charge of soils investigations for the duration of the project. Field research involved close collaboration with the Soil Survey Laboratory staff, particularly the Lincoln, Nebraska unit headed by R. B. Grossman. Work was done cooperatively with the New Mexico State University College of Agriculture, and Departments of Biology and Earth Sciences, and the New Mexico Bureau of Mines and Mineral Resources. The NMSU Agricultural Experiment Station and Department of Agronomy provided office and laboratory space, and numerous other supporting services for the 15-year period [*cf.* Monger et al. 2009].

The project area is typical of large parts of west-central North America, and results of investigations can be applied in many other warm-arid regions, particularly those with basin and range topography [**Fig. C-14**]. The area comprises the northern Mesilla Valley segment of the Rio Grande, parts of two internally-drained intermontane basins [Jornada del Muerto and Mesilla], and several mountain masses. Because of striking differences in rock units exposed in source watersheds in local mountains and the upper Rio Grande basin (e.g. rhyolite, monzonite, limestone, and mixed-[composition ARG] fluvial deposits), it has been possible to study effects of various bedrock and alluvial parent materials on desert soil development [Gile 1961; Gile and Grossman 1979; Gile et al. 1965, 1966; 1981].

Field investigations in the project area included mapping of geomorphic surfaces, geologic units, and soils at scales ranging from 1:8,000 to 1:24,000. Detailed studies were also made in the Rincon Valley area of northern Dona Ana County [Gile et al. 1996], and reconnaissance investigations were conducted at a number of sites elsewhere in the Basin and Range province. Geologic studies emphasized late Cenozoic stratigraphy, geomorphic surface mapping, and sedimentology and hydrogeology of basin and valley fills [e.g., Hawley and Gile 1966; Hawley 1969b, 1970, 1972, 1975a and b; Hawley and Kottlowski 1969; Hawley et al. 1969; King et al. 1969, 1971; Seager and Hawley 1973; Seager et al. 1971, 1975; King and Hawley 1975]. Pedologic

**Recipient of the 1983 Geological Society of America - Kirk Bryan Award: for paper of distinction advancing the science of geomorphology (cf. Birkeland 1984).*



C3.2. Research Collaborations Related to the NM Water Research Institute Program

Early hydrogeologic phases of Desert Project studies involved collaboration with the USGS-WRD in locating water-supply wells for the new NASA-Apollo Project White Sands Test Facility (Doty 1963). Formal collaboration with water-resources research programs at NMSU was initiated **when** Hawley was invited to be a Charter Member of the newly established NM Water Resources Research Institute by Director, Dr. H.R. Stucky, in October 1964 (Fig. C3-2).




Figure C3-2. Facsimile copy of letter of 10/15/1964 from NM WRRI Director H.R. Stucky to John Hawley concerning the latter's appointment as an Institute "Charter Member."

Initiation of NM Water Resources Research Institute programs in 1964 reinforced the pressing need for a fully accredited geology department at NMSU. In this respect, one of Dr. Ralph Stucky's earliest contributions to groundwater science in the southern New Mexico area was the result of his forceful advocacy with the NMSU Board of Regents for formation of an undergraduate-major program in the combined earth-science fields of geology and geography. This activity's early success was exemplified in the 1964 appointment of Dr. William E. (Bill) King (1925-2014, a former Mobil-Oil Senior Geologist) as Chairman of a new NMSU Earth Science & Astronomy Department. The following excerpt from the introduction to an early NM WRRI report (No. 6) on the "Hydrogeology of the Rio Grande Valley and adjacent intermontane areas of southern New Mexico" illustrates the substantial contributions to the understanding of the Mesilla Basin region in the first five years of NM WRRI's existence (King et al. 1969, p. 5):

In 1964, the Water Resources Research Act was passed by the United States Congress, and the Water Resources Research Institute of New Mexico, under the direction of H. R. Stucky, was established soon thereafter. The present investigation was among the first to be funded by the institute. The field and laboratory work was begun in February 1965 and terminated in August 1968.

The principal investigator, W. E. King, professor of geology at New Mexico State University, spent one quarter of his time through each academic year and three full months each summer on the investigation. He was responsible for supervision of the investigation and is accountable for many of the conclusions. Andrew M. Taylor [Taylor 1967] served as a graduate assistant on the project from its inception through August 1967 and did much of the well logging and surface geology. Richard P. Wilson spent one year as a student assistant and is largely responsible for the drafting as well as much of the thought expressed in the water-table contour map (Pl. 1).

John W. Hawley, areal geologist with the Soil Conservation Service, of the U.S. Department of Agriculture, suggested the investigation and cooperated in the study throughout [cf. King and others (1971) p. 9-24].*

***Part C8a** is a special Appendix Addendum that comprises a reformatted copy of the section by J. W. Hawley on the "Geology and its relation to the hydrologic system" in King, W.E. and others, 1971, *Geology and ground-water resources of central and western Doña Ana County, New Mexico: New Mexico Bureau of Mines and Mineral Resources, Hydrologic Report 1*, p. 9-24.

C4. USGS-WATER RESOURCES DIVISION INVESTIGATIONS IN THE DOÑA ANA COUNTY, NM AREA: THE CLYDE A. WILSON ERA—1971 to 1980.

C4.1. Background

Reports by Clyde A. Wilson and USGS Water-Resources Division (WRD) associates, which were published by the NM OSE and the USGS, remain the most comprehensive accounts of geohydrological investigations in the area of Mesilla Basin and Rincon Valley/Palomas Basin between Caballo Dam and the International Boundary (**Fig. C4-1**; Wilson et al. 1981, Wilson and White 1984; cf. Gates and Stanley 1976 [Report **Part 3.7.3, Fig. 3-16**]). Due to Wilson's unanticipated death in October 1980, only a fraction of his impressive geohydrologic and geophysical work was ever published. However, with full support of the USGS-WRD New Mexico and Texas Districts, Hawley and NM Tech colleagues were able to acquire a nearly complete set of Wilson's detailed, handwritten notes on a large number of deep test-well installations, many of which were supplemented by borehole-sample and geophysical logs. This subsurface database has been of immeasurable value in all subsequent hydrogeologic investigations in the Mesilla Basin region (e.g., Khaleel et al. 1983, Peterson et al. 1984, Hawley and Lozinsky 1992, Nickerson and Myers 1993, Hawley and Kennedy 2004, and Hawley et al. 2005). Additional information on Wilson's seminal contributions is included in **Parts C4.3** and **4.4**.

PROJECT PROPOSAL* FOR A COMPREHENSIVE STUDY OF THE WATER RESOURCES
OF THE SOUTHERN RIO GRANDE VALLEY AREA, NEW MEXICO

C. A. Wilson

UNITED STATES DEPARTMENT OF THE INTERIOR, GEOLOGICAL SURVEY
Albuquerque, New Mexico, May 1972

** **Part C8b** is a special Appendix Addendum that comprises a reformatted copy of the heretofore unpublished "Project Proposal" by C.A. Wilson (5/1972).*

INTRODUCTION [p. 2-3]

The water supply of the lower Rio Grande Valley and adjacent uplands is derived from surface water from the Rio Grande and ground water from sediments of the intermontane basin. Neglecting the short-term effects, the surface and ground-water units function as one system, due to the hydraulic connection between the river and the aquifers. Planning the orderly development of water resources of the valley requires current factual data followed by complete and reliable analyses.

The purpose of this proposal is to aid in planning, organization, and preparation of a comprehensive study plan of the southern Rio Grande Valley in New Mexico, to evaluate data presently available, and to determine data that will need to be collected (*cf.* **Fig. C4-2**).

CONTENTS

INTRODUCTION	4
NEED FOR THE STUDY	8
PREVIOUS STUDIES AND DATA AVAILABILITY.....	13
OBJECTIVES AND PLAN OF STUDY	18
Water resources in vicinity of Las Cruces.....	22
Water resources in the Rincon and Mesilla Valleys.....	24
Major items of work.....	30
Schedule of work, time, and costs.....	34
Reports.....	40
SELECTED REFERENCES	42

ILLUSTRATIONS

Figure 1 (<i>cf.</i> Fig. C4-1). – Map of lower Rio Grande Valley area, New Mexico	5
---	---




Figure C4-1. Facsimile copy of Figure 1. "Location of study area" from Wilson and others (1981, p. 5).



United States Department of the Interior

GEOLOGICAL SURVEY
Water Resources Division
P.O. Box 4198
El Paso, Texas 79914

August 14, 1972

Dr. John W. Hawley
U. S. Soil Conservation Service
1616 19th St.
Lubbock, Texas 79403

Dear John:

As you probably know from talking to Clyde Wilson and Jim Smith, we are beginning an evaluation of ground water in several areas in West Texas. One area we will concentrate on is the Mesilla Valley, where we plan to run several earth-resistivity profiles this fall and drill 2-3 deep test holes next year in an attempt to better define the potential of the Santa Fe aquifers. I am rounding up references on the Cenozoic geology of the area, and would like to obtain a copy of your 1966 Friends of the Pleistocene guidebook. Could you tell me if they are still available and how to get one?

You probably won't recall, but we met briefly on a Friends of the Pleistocene trip at Promontory Point, Utah in 1958 or 1959. The trip was to look at the Lake Bonneville deposits exposed by excavation for the SP causeway. As I recall you came over from Nevada with George Maxey. Sometime when you're in Las Cruces, we would appreciate your taking a day and showing us some of the pertinent features of the Mesilla Valley.

Sincerely

Joseph S. Gates
- J.S.G.
Joseph S. Gates
Hydrologist

JSG: bf

Figure C4-2. Facsimile copy of 8/14/1972 letter from Joseph S. Gates (USGS-WRD) on Proposed Earth-Resistivity Surveys and Test Drilling in the El Paso area of the Southern Mesilla Basin (*cf.* Gates and Stanley 1976, Jackson 1976, Zohdy et al. 1976, Wilson et al. 1981).

C4.2. Major Contributions by C.A. Wilson to the Understanding of Regional Hydrogeology in Water Resources of the Rincon and Mesilla Valleys and Adjacent Areas in New Mexico: New Mexico State Engineer Technical Report 43 (Wilson et al. 1981)

The scope of investigations directed by C.A. Wilson and colleagues and their major contributions to the current understanding of the basic hydrogeologic-framework components in the Mesilla Basin region is illustrated in the following selections from parts of the "Methods of investigation" section in Wilson and others (1981, p. 13-16):

1. Well inventory

During the study, about 1,530 water wells were inventoried for such information as total depth, static and pumping depth-to-water, yield, and the specific conductance and temperature of the pumped water. . . .

2. Test drilling

Test holes ranging in depth from 135 to 2,470 feet were drilled at 27 locations in the study area. Hole diameters ranged from about 6 to 8 inches. Drill cutting samples from each test hole were used to compile a geologic log. This log was later revised on the basis of microscopic examination of cuttings and examination of available geophysical logs. Borehole geophysical logs, including natural gamma-ray, spontaneous potential, and resistivity logs, were made in 25 of the holes. Neutron, gamma-gamma (density), and caliper logs also were recorded in selected deep holes. . . .

3. Geophysical studies

Geophysical studies, which included surface electrical-resistivity soundings, permitted the estimation of critical hydrologic data from depths below the limits of existing wells. Vertical electrical-resistivity soundings (VES) made on the land surface permitted estimates of subsurface rock properties and ground-water salinity and allowed coverage of large areas in a short time and at a reasonable cost. . . .

Electrical-resistivity sounding were made at 150 sites in the Rincon and Mesilla Valleys and adjacent areas (plates 1 and 2*). The initial set of 50 soundings was made during 1973 by Geoterrex, Ltd., Tucson, Arizona. The remaining 100 sounding were made by the U.S. Geological Survey (Jackson, 1976). Sites were selected so that results from the soundings could be integrated with borehole logs and chemical analyses of water from test holes and water wells in order to improve the geophysical log interpretation. . . . [cf. Jackson 1976, Jackson and Bisdorf 1975, Zohdy et al. 1976].*

**Note: VES-survey and borehole-log hydrogeologic interpretations are illustrated in 14 "Hydrologic sections" of basin-fill deposits in the Rincon and Mesilla Valley areas (respectively, A-A' and B-B' and C-C' to N-N'). Section-base elevation is 4,000 ft (1,219 m) below msl (Wilson et al. 1981, plates 1, 2, and 8).*

With regards to current estimates of the amount of fresh and slightly saline groundwater stored in basin-fill aquifers of the Mesilla Basin's RG Valley and "West Mesa" area, the following selections from Wilson and others (1981, p. 58-59) are of special relevance:

The approximate thickness of sediments saturated with freshwater ranges from less than 400 feet in the northern and southern ends of the valley to more than 2,400 feet in the central valley (plate 15). The thickest part of the freshwater zone generally follows the present course of the Rio Grande.

In the part of the Mesilla Valley north of Anthony, the volume of sediments containing freshwater is estimated to be approximately 215 million acre-feet. If the average sediment porosity is assumed to be 30 percent, about 64 million acre-feet of freshwater are in storage. Assuming that all recharge to the aquifer stops, that pumpage continues for a very long time, that average effective porosity is 15 percent, and that 60 percent of the sediments are sand and gravel, approximately 20 million acre-feet of fresh-water may be available for withdrawal by wells. Any recharge from applied irrigation water, river or canal seepage, inflow from other areas, precipitation, and water that moves from clay units into sands in the aquifer increases the volume of freshwater available for pumping.

In the part of the Mesilla Valley south of Anthony, much less freshwater is present. Leggat, Lowry, and Hood (1962, p. 38-39) estimated that about 980,000 acre-feet of freshwater are in storage in the basin fill of the New Mexico part of the Mesilla Valley that is south of Anthony and

east of the western edge of the flood-plain alluvium. They defined freshwater as water that contains less than 250 parts per million chloride. Gates, White, Stanley, and Ackermann (1978 [1980]) prepared a map of the Mesilla Valley south of Anthony showing the approximate thickness of basin fill containing freshwater. They (1978 [1980], p. 108) estimated that 820,000 acre-feet of freshwater are in storage beneath the Texas part of the Mesilla Valley and the adjacent mesa to the east.

Adding the estimates of freshwater in storage north of Anthony (64 million acre-feet) and south of Anthony (1.8 million acre-feet) gives a total of about 66 million acre-feet. The saturated thickness of sediments containing freshwater beneath the West Mesa ranges from less than 400 to more than 2,400 feet (plate 15). The greater thickness occurs near the Mesilla Valley.

The volume of saturated sediments in the freshwater zone beneath the West Mesa is estimated to be 380 million acre-feet. Assuming an average porosity of 30 percent, about 114 million acre-feet of freshwater may be in storage. Not all of this water would be available to wells. Assuming that the specific yield of the sand layers is 15 percent and that 60 percent of the freshwater zone is sand, about 34 million acre-feet of freshwater theoretically could be pumped in the West Mesa. Additional water may be derived from the clay units in the aquifer. The very large volume of slightly saline water [SSL] that lies beneath the freshwater zone is not considered in these calculations although this water may be suitable for some purposes. . . .

The upper SSW zone is thinnest on the west side of the Mesilla Valley and thickens to the east where it unites with the slightly saline water that underlies the freshwater zone (plate 8, sections C-C' through N-N'). In the Mesilla Valley area extending south to Anthony, the volume of sediments containing slightly saline water that lies above the zone of freshwater is estimated to be 17 million acre-feet. Assuming a porosity of 30 percent, there are about 5.1 million acre-feet of slightly saline water in storage. The upper SSW zone is generally under unconfined conditions. Assuming 80 percent of the zone is sand and gravel and the average specific yield of the sand and gravel is 20 percent, about 2.7 million acre-feet of slightly saline water theoretically would be available to shallow wells in the Mesilla Valley north of Anthony. This upper SSW zone furnishes much of the water to the valley's irrigation wells.

C4.3. Memorials to Clyde A. Wilson (1932-1980)

C4.3.1. Memorial to Clyde Wilson *in* Wilson and others (1981, p. ii)

Clyde A. Wilson, the senior author of this report, died on October 11, 1980, in the Gila National Forest in southern New Mexico. Only through his guidance and expertise was this study possible.

C4.3.2. Memorial to Clyde Wilson in the NM WRRI *The Divining Rod*:

Clyde Wilson Dies at Age 48 (Mackichan, 1980, *The Divining Rod*, no. 80-3. p. 1, 3):

Clyde Wilson, U.S. Geological Survey hydrologist at Las Cruces, New Mexico, died Saturday, October 11, 1980, while on a family hunting trip. He had been chief of the Las Cruces field office since 1971. Clyde was born in Morenci, Arizona in 1932. After graduating from Carlsbad High School in 1950, he first attended New Mexico School of Mines, Socorro. He then enlisted in the Air Force and received his commission and wings in March 1954. Wilson served as a navigator-bombardier in Korea, and an Air Force police officer at Hurlbut Field, Florida until March, 1957, when he left the service with the rank of First Lieutenant.

Returning to college, Clyde received his B.S. in geology from Texas Technological College, Lubbock, and his M.S. in geology from the University of Arizona. He later earned a second Master's degree in hydrology from the University of Arizona. Wilson had been employed by the U.S. Geological Survey since 1962. He had worked as a geologist on several Texas projects, and later was assigned to the Permian Basin Project in Austin, Texas, as a hydrologist. Clyde came to Las Cruces from Lubbock, Texas, where he was involved in a study of the Ogallala Aquifer.

When Clyde first came to Las Cruces, he developed a proposal for a groundwater study of the lower Rio Grande [NM] area [Part C7.2]. In July, 1972, the formal study began with the Office of State Engineer . . . , the City of Las Cruces, and the Bureau of Reclamation for the Elephant Butte Irrigation District as cooperators. It was at this time that the USGS moved its office into Stucky Hall with the WRRI on the NMSU campus. . . ,

C5. MULTI-INSTITUTIONAL COOPERATIVE INVESTIGATIONS—1980 to 2000

C5.1. Beginning of the “Administratively Closed Groundwater-Basin” Era

C5.1.1. Management of Shared-Water Resources—Progress and Pitfalls

The following excerpt from Harris (2012, p. 249) serves as a very appropriate introduction to the following discussion:

On Sept. 5, 1980, El Paso took the first step along a legal route it hoped would lead to a plentiful and free water supply from New Mexico. On that Friday after Labor Day, the city of El Paso, through the Public Service Board, filed suit in US. District Court in Albuquerque against Steve Reynolds individually as the New Mexico state engineer, and also New Mexico Attorney General Jeff Bingaman and the New Mexico district attorney for Doña Ana County. The city sought to overturn New Mexico's embargo statute as violating the Commerce Clause of the US. Constitution. The embargo statute, enacted in 1953, prevents anyone from drilling wells in New Mexico and transporting the water outside the state.

By September 12, New Mexico State Engineer Steve Reynolds "declared" both the Mesilla Basin and the Hueco Bolson under state authority. El Paso responded quickly by filing well applications for 246,000 acre-feet a year in the Mesilla Basin* and for 50,000 acre-feet a year in the Hueco Bolson.

**USGS hydrologist Clyde A. Wilson, the leading expert on Mesilla Basin and Lower Rio Grande Valley water-resource matters, died unexpectedly on October 11, 1980*

The state engineer denied all of El Paso's applications on grounds that New Mexico statute prohibited the transfer of water across the state line. El Paso took its case before US. District Court Judge Howard Bratton in January 1982. In July of that year the US. Supreme Court ruled that groundwater was to be considered an article of commerce and as such its transfer across state lines could not be restricted. In January 1983, Bratton ruled in favor of El Paso on all accounts, writing that New Mexico's embargo violated the Commerce Clause of the US. Constitution because it promoted New Mexico's economic advantage.

C5.1.2. Hydrogeologic Framework Characterization for Numerical Modeling

Hydrogeologic investigations between 1983 and 1997 included collaborative efforts involving the NMBMMR and NM Tech (Hawley and Rick Lozinsky), USGS-WRD (Mike Kernodle and Ken Stevens), NMOSE (Francis West) and EPWU (Tom Cliett [Part C-2b]). Emphasis was on compilation and interpretation of subsurface geologic, geophysical and geochemical data (e.g., Hawley and Lozinsky 1992, Nickerson and Myers 1993, Cliett and Hawley 1996). The focus of most hydrogeologic-research activities was on development of framework characterizations of basin-fill aquifer systems of the Mesilla Basin region that were compatible with emerging numerical groundwater-flow modeling, computer-platforms, and GIS technologies.

Much of the basic information compiled for studies initiated after 1982 had already been collected during early stages of geohydrologic and hydrogeologic research projects jointly sponsored by the USGS-WRD, NMBMMR, NM Mexico Tech Research Division, NM WRRI, and NMSU-Energy Institute (Parts C2 to C4). In addition to published sources, available unpublished, subsurface data (primarily drilling, and borehole-sample and geophysical logs) was obtained from USGS-WRD and

NMBMMR files. Emphasis was on development of a lithofacies classification scheme for use in precursor (quasi-3D) groundwater-flow models of basin-fill (alluvial) aquifer systems (e.g., Khaleel et al. 1983, Hawley 1984, Peterson et al. 1984; *cf.* **APNDX A**).

Published-information sources utilized in those studies came from three complementary earth-science disciplines:

1. “Traditional” field geology and geophysics exemplified by work synthesized by Hawley and others (1969), Hawley (1975a, 1978), Seager and Morgan (1979), Seager (1981), Gross and Icerman (1983), Icerman and Lohse (1983), Seager and others (1987), Keller and others (1990), Seager and Mack (1994), Mack and others (1998), and Seager (1995). *See* **CHAPTER 3** and **APPENDIX H**.
2. Integrated soil-science and Quaternary geology/geomorphology studies, notably work by Gile and Hawley (1966 and 1968), Gile and Grossman (1979), and Gile and others (1981, 1995). *See* Monger and others (2009) and **APPENDIX D**.
3. Integrated geohydrologic and hydrogeologic studies by Conover (1954), Knowles and Kennedy (1958), Leggat and others (1962), King and others (1971), Wilson and others (1981), Wilson and White (1984), Frenzel and Kaehler (1992), and Nickerson and Myers (1993).

Key sources of borehole data were identified and located on available geologic maps of the Mesilla Basin (scales 1:24,000 and 1:100,000) for use as control points. These included borehole geophysical and sample logs, geothermal data, and geochemical analyses. Six new test wells drilled by the USGS-WRD and EPWU provided supplemental information. The Afton, Lanark, La Union, and Noria test wells were drilled in the basin area west of Mesilla Valley (MT 1 to 4; Well nos. 175, 236, 242, and 253 *in* Report **TABLE 1**). The other two wells (CWF1D, CWF4D; JL-49-04-481 and 469; Report **TABLE 1** nos. 343 and 341) are located in the Canutillo Well Field area on the Rio Grande floodplain west of Vinton, Texas (Frenzel and Kaehler 1992, Hawley and Lozinsky 1992; Nickerson 1989, 1995; Nickerson and Myers 1993).

Drill-cutting analysis and driller-log data (Report **TABLE 1**) were supplemented by detailed seismic reflection profiles made at two sites near the Canutillo Well Field (C.B. Reynolds and Associates 1986, 1987 *unpublished*). The subsurface database also included water-chemistry analyses from one or more sampling intervals in most of the key wells (Hawley and Lozinsky 1992, Table 4; Appendix—Tables A1, A2, A3).

C5.2. USGS-WRD Regional Aquifer-Systems Analysis (RASA): Southwest Alluvial Basins (SWAB) Program

The USGS-WRD Regional Aquifer-Systems Analysis (RASA)—Southwest Alluvial Basins (SWAB) Program has had a major impact on development of new-generation conceptual and digital models of intermontane-basin hydrogeologic systems throughout the eastern Basin and Range and Rio Grande-rift provinces (*cf.* Gates et al. 1984, Freethy et al. 1986). The following selections from Frenzel and Kaehler (1992, p. iii), and (Wilkins 1986, p. 2) summarize the program’s purpose and scope:

THE REGIONAL AQUIFER-SYSTEM ANALYSIS PROGRAM

The Regional Aquifer-System Analysis (RASA) program was started in 1978 after a congressional mandate to develop quantitative appraisals of the major ground-water systems of the United States. The RASA program represents a systematic effort to study a number of the Nation’s most important aquifer systems that, in aggregate, underlie much of the country and that represent important components of the Nation’s total water supply. In general, the boundaries of these studies are identified by the hydrologic extent of each system and accordingly transcend the political subdivisions to which investigations have often arbitrarily

been limited in the past. The broad objective for each study is to assemble geologic, hydrologic, and geochemical information; to analyze and develop an understanding of the system; and to develop predictive capabilities that will contribute to the effective management of the system. The use of computer simulation is an important element of the RASA studies, both to develop an understanding of the natural, undisturbed hydrologic system and any changes brought about by human activities as well as to provide a means of predicting the regional effects of future pumping or other stresses. . . .

The objectives of Regional Aquifer-Systems Analysis (RASA) are: (1) to describe the water-resource system, (2) to analyze changes in the system, (3) to develop a data base from existing information, and (4) to simulate the hydrologic system using mathematical models.

Hydrologic data, geologic descriptions of basin boundaries, and aquifer properties presented in this report were compiled from the many previous studies completed in the regional study area. Regional interpretations resulting from analysis of existing data are also presented in this report.

The study area encompasses parts of Colorado, New Mexico, and Texas and is structurally and hydrologically divided into 22 surface-water open and closed basins [Fig. C5-1]. The Rio Grande is the primary hydrologic connection through the open basins of southern Colorado, New Mexico, and western Texas. Closed basins in southwestern New Mexico and western Texas make up the rest of the area. . . .




Figure C5-1. Location of the Mesilla Basin (12) within the RASA “Southwest Alluvial Basin Regional Aquifer-System Analysis (SWAB)” area of the United States (from Frenzel and Kaehler 1992, Fig. 1; *cf.* Wilkins 1986, 1998). Major basins of the Rio Grande rift (RG-rift tectonic province: 1-10, east 14, and 19-22; with other contiguous Basin and Range physiographic-province basins: 14-18). Besides the Mesilla Basins, the Study Area includes parts of the following groundwater basins: Southern Jornada, and western Tularosa-Hueco (13).

In addition to Frenzel and Kaehler (1990, 1992), the following reports are representative of the substantial contribution made by SWAB-Program investigations to NM WRRI-sponsored hydrogeologic research in the Mesilla Basin region:

- Anderholm, S.K., 1985, Clay-size fraction and powdered whole-rock X-ray analyses of alluvial basin deposits in central and southern New Mexico: U.S. Geological Survey Open-File Report 85-163, 18 p.
- Freethy, G.W., Pool, D.R., Anderson, T.W., and Tucci, P., 1986, Description and generalized description of aquifer materials in alluvial basin of Arizona and adjacent parts of California and New Mexico: U.S. Geological Survey Hydrologic Investigations Atlas HA-664, 4 sheets, scale 1:500,000.
- Hanson, R.T., McLean, J.S., and Miller, R.S., 1994, Hydrogeologic framework and preliminary simulation of ground-water flow in the Mimbres Basin, southwestern New Mexico: U.S. Geological Survey Water-Resources Investigations Report 94-4011, 118 p.
- Kernodle, J.M., 1992a, Results of simulations by a preliminary numerical model of land subsidence in the El Paso, Texas, area: U.S. Geological Survey Water-Resources Investigations Report 92-4037, 35 p.
- Kernodle, J.M., 1992b, Summary of U.S. Geological Survey ground-water-flow models of basin-fill aquifers in the southwestern alluvial basins region, Colorado, New Mexico, and Texas: U.S. Geological Survey Open-File Report 90-361, 81 p.
- Myers, R.G., and Orr, B.R., 1986, Geohydrology of the aquifer in the Santa Fe Group, northern West Mesa of the Mesilla Basin near Las Cruces, New Mexico: U.S. Geological Survey Water-Resources Investigations Report 84-4190, 37 p.
- Nickerson, E.L., and Myers, R.G., 1993, Geohydrology of the Mesilla ground-water basin, Doña Ana County, New Mexico, and El Paso County, Texas: U.S. Geological Survey Water-Resources Investigations Report 92-4156, 89 p.
- Orr, B.R., and Risser, D.W., 1992, Geohydrology and potential effects of development of freshwater resources in the northern part of the Hueco Bolson, Doña Ana and Otero Counties, New Mexico, and El Paso County, Texas: U.S. Geological Survey Water-Resources Investigations Report 91-4082, 92 p.

A significant contribution of the first major RASA-SWAB Project completed in New Mexico (Frenzel and Kaehler 1990, 1992) involved making a formal distinction between the deep, structural, Mesilla “groundwater basin (MeB),” and the areally more-extensive, physiographic- and hydrographic-basin categories, with the latter being defined, respectively, in terms of Basin and Range topography and surface-watershed divides (**Figure C5-2**). The MeB boundaries selected by USGS hydrologists Peter Frenzel and Charles Kaehler were specifically designed for an initial-phase “digital model of the Mesilla Basin ground-water flow system (1992, p. C2, Fig. 4 [1990, p. 9-10]).” Except for its southern extension into Chihuahua, these boundaries closely match mapped positions of the Study Area’s major Basin and Range/RG-rift basin-border fault zones (e.g., L. Woodward et al. 1978, Seager et al. 1987, and Seager 1995). Accordingly, MeB boundary locations in all subsequent digital hydrogeologic-framework characterizations and derivative GW flow-models in the USA part of Mesilla Basin region remain in close agreement with those defined by Frenzel and Kaehler (1990, 1992; Report **PLATES 1 to 5**; cf. Hawley and Lozinsky 1992, Nickerson and Myers, 1993, Hamilton and Maddock 1993, Hawley and Kennedy 2004, Sweetkind 2017).




Figure C5-2. Map showing locations of the Mesilla Basin, Mesilla Valley, and bounding bedrock uplifts in the Doña Ana County area of south-central New Mexico. Dashed line shows boundary of the Mesilla “groundwater basin [MeB]” as defined by P.F. Frenzel and C.A. Kaehler (1990, Fig. 4; 1992, Fig. 4).

While none of the above-cited reports included information on potentiometric-surface elevation or regional groundwater-flow direction south of the International Boundary, invaluable interpretations on approximate predevelopment water levels are presented in maps compiled by Conover (1954), King and others (1971), Wilson and others (1981), and Nickerson and Myers (1993) for the 1946 to 1985 interval (*see* Report **CHPT. 7**). All, however, show a regional GW-flow component that is subparallel to binational-border zone with a focused-outflow discharge area in the southernmost Mesilla Valley area between Canutillo and Sunland Park (*cf.* **Parts C2.1 and C2.2, Figs. C2-1 and C5-3**). Moreover, most also indicate that a substantial flow component had a recharge source in north-central Chihuahua's "Zona Hidrogeológica de Conejos Médanos" (Conover 1954, Plate. 1 [**PLATE 9A**]; King et al. 1971, Plate. 1 [**PLATE 9B**]; Wilson et al. 1981, Pl. 9; Nickerson and Myers 1993, Fig. 9). **Figure C5-3** (south part of **PLATE 9C**) shows the approximate pre-development water-level contours in the southwestern Mesilla Basin. **Figure C5-4** (from **PLATE 4**) is included to illustrate the remaining degree of uncertainty on the nature of the GW-flow regime in parts of the Study Area in Chihuahua.




Figure C5-3. (Plate 9C extract) Map showing approximate water-level contours and gaging stations on drains, Mesilla Basin, New Mexico, and Texas (Lower part of Plate 1 in Frenzel and Kaehler 1992 [**Plate 9C**]; modified from Wilson et al. 1981 [Plate 9]—1976 water-level compilation) (*cf.* **Figure C5-4** and **PL. 4**).




Figure C5-4. (page-size **PLATE 4**) Index map of major hydrographic and geohydrologic features of the Study Area. Surface-watershed boundaries—heavy-blue lines. Pre-development potentiometric-surface elevation approximation (10-ft contour interval) in central-basin areas—light-blue lines. The dashed-blue in the map’s SW corner shows the approximate position of the GW-flow divide between lower Mesilla Valley directed flow and Los Muertos Basin/El Barreal-directed flow. Swanson Geoscience, LLC compilation on a **PLATE 1** hydrogeologic map base (1:100,000 compilation scale).

C5.3. USGS-State Cooperative Studies of Geology and Hydrology in the Basin and Range Province Related to Selection of Sites for the Disposal of High-Level Radioactive Waste

Throughout the 1980s, USGS-State collaborative studies of the Basin and Range province relative to the disposal of high-level radioactive waste also had a significant impact on the development of present generation of hydrogeologic-framework models of the Mesilla Basin region, particularly with respect to groundwater-flow regimes in interlinked rift-basin interiors and inter-basin bedrock uplifts (Bedinger et al. 1984, 1989a-c). The following selections from the “INTRODUCTION” to USGS Professional Paper 1370-A describe the program’s “background and purpose” (Bedinger, Sargent, and four others, 1989, p. A-2):

A study by the U.S. Geological Survey to evaluate potential hydrogeologic environments for isolation of high-level radioactive waste in the Basin and Range physiographic province was begun in May 1981, with the introduction of the study to the Governors of eight Basin and Range States-Arizona, California, Idaho, Nevada, New Mexico, Oregon, Texas, and Utah-and to Indian tribes in those States. Accordingly, these States were invited to participate in the study by designating an earth scientist to serve on a Province Working Group with the U.S. Geological Survey-membership of the working group is shown following the title page. State representatives have provided consultation in selecting guidelines, assembling geologic and hydrologic data, and assessing such information to identify environments that meet the guidelines for further study....

The regions were selected [for further evaluation during first-phase studies] on the basis of the adopted guidelines and of information obtained from available sources on: (1) The distribution of rock types that may be host media for radioactive waste; (2) characteristics of the province related to tectonic stability-seismicity, late Cenozoic volcanism, Quaternary faulting, late Cenozoic regional uplift, and heat flow; and (3) the hydrology of ground-water flow systems.

This report, Chapter A of Professional Paper 1370, is one of a series of eight chapters (A through H) that describe the evaluation of potential hydrogeologic environments for the isolation of high-level radioactive waste in the Basin and Range physiographic province. These chapters present the results of the second phase of study. The titles of chapters in this series are:

- A Basis of characterization and evaluation
- B Characterization of the Trans-Pecos region, Texas
- C Characterization of the Rio Grande region, New Mexico and Texas
- D Characterization of the Sonoran region, Arizona
- E Characterization of the Sonoran region, California
- F Characterization of the Death Valley region, Nevada and California
- G Characterization of the Bonneville region, Utah and Nevada
- H Evaluation of the regions.

. . . Resources and time did not permit study in the present, second, phase of all potential regions identified in the first phase of study. . . The regions selected for study and revised subdivisions are shown in figure 2 (**Fig. C5-5**)




Figure C5-5. (Bedinger et al. 1989a, Fig. 2) Map of regions (shaded) in the Basin and Range physiographic province selected for evaluation of “potential hydrogeologic environments for isolation of high-level radioactive waste.” “Rio Grande Region (RG):” Tularosa Basin and Hueco Bolson-01, and Central and Northern Jornada (del Muerto) Basin-02 (Bedinger et al. 1989b, Johnson et al. 1989, and Sargent et al. 1989).

C6. CONCLUDING REMARKS

The following selection from Nickerson (2006, p. 2) represents the first formal USGS recognition of the hydrogeologic-framework model that was developed by Hawley and Lozinsky (1992) for use throughout the Mesilla Basin region (*cf.* Hawley et al. 2009, Sweetkind 2017, Sweetkind et al. 2017, Teeple 2017a, Hanson et al. 2018):

The lower Mesilla Valley ground-water system consists primarily of a basin-fill aquifer with two main geologic units: the Santa Fe Group of Quaternary and Tertiary age and the Rio Grande flood-plain alluvium of Quaternary age (King and others, 1971). The Santa Fe Group is an intermontane basin-fill unit that extends throughout the Mesilla Basin and includes alluvial, eolian, and lacustrine deposits (Hawley and Lozinsky, 1992, p. 4). These clay, silt, sand, and gravel deposits can reach depths of more than 2,000 feet in the lower Mesilla Valley. The Rio Grande flood-plain alluvium overlies the Santa Fe Group in the Mesilla Valley and consists of channel and flood-plain deposits of clay, silt, sand, and gravel that generally are less than 125 feet thick (Wilson and others, 1981, p. 27).

The hydrogeologic framework of the Mesilla Basin was established in Hawley and Lozinsky (1992). Their basinwide conceptual model delineated three hydrogeologic features: (1) bedrock and structural boundaries, (2) lithofacies assemblages, and (3) hydrostratigraphic units. Santa Fe Group basin fill is divided into the informal upper, middle, and lower hydrostratigraphic units based on depositional environment and age. The upper Santa Fe hydrostratigraphic unit (USF) consists primarily of ancestral Rio Grande deposits of medium to coarse sand and [pebble] gravel. The USF includes the Camp Rice and upper Fort Hancock Formations. The middle Santa Fe hydrostratigraphic unit (MSF) consists primarily of alluvial deposits with eolian and playa-lake facies.

Basin-floor sediments of interbedded sand and silty clay are common. The MSF includes the Fort Hancock and Rincon Valley Formations. The lower Santa Fe hydrostratigraphic unit (LSF) consists primarily of eolian, playa-lake, and alluvial facies. Basin-floor sediments include thick-bedded dune sand. The LSF includes the Hayner Ranch and the lower Rincon Valley Formations. Detailed descriptions of the hydrostratigraphic units and associated lithofacies can be found in Hawley and Lozinsky (1992). The Mesilla Basin hydrogeologic framework was recently updated and integrated into a digital format by Hawley and Kennedy (2004).

C7. CITED REFERENCES AND NOTABLE CONTRIBUTIONS

C7.1. Cited References (with Topical/Sub-Topical Alphanumeric Cross-Reference Codes, see Appendix B)

- Allen, B.D., 2005, Ice Age lakes in New Mexico, *in* Lucas, S.G., Morgan, G.A., and Zeigler, K.E., eds., *New Mexico's Ice Ages: New Mexico Museum of Natural History and Science Bulletin* 18, 107-114. **(C1, I2)**
- Anderholm, S.K., 1985, Clay-size fraction and powdered whole-rock X-ray analyses of alluvial basin deposits in central and southern New Mexico: U.S. Geological Survey Open-File Report 85-163, 18 p. **(C2a, H2)**
- Anderson, T.W., 1995, Summary of the southwest alluvial basins, regional aquifer-system analysis, south-central Arizona and parts of adjacent states: U.S. Geological Survey Professional Paper 1406-A, 33 p. **(D1, F2)**
- Anderson, T.W., Welder, G.E., Lesser, G., and Trujillo, A., 1988, Region 7, Central alluvial basins, *in* Back, W., Rosenshein, J.S., and Seaber, P.R., eds. *Hydrogeology—The Geology of North America: Geological Society of America, DNAG Volume. O-2*, p. 81-86. **(D1)**
- Barker, F.C., 1898, Irrigation in Mesilla Valley, New Mexico: U.S. Geological Survey Water-Supply and Irrigation Paper 10, 51p. **(G1)**
- Bedinger, M.S., Sargent, K.A., and Langer, W.H., eds., 1989a, Studies of geology and hydrology in the Basin and Range province, southwestern United States, for isolation of high-level radioactive waste – Characterization of the Trans-Pecos region, Texas, U.S. Geological Survey Professional Paper 1370-B, 43 p. **(F2, H3)**

- Bedinger, M.S., Sargent, K.A., and Langer, W.H., eds., 1989b, Studies of geology and hydrology in the Basin and Range province, southwestern United States, for isolation of high-level radioactive waste – Characterization of the Rio Grande region, New Mexico and Texas: U.S. Geological Survey Professional Paper 1370-C, 48 p. **(F2, H3)**
- Bedinger, M.S., Sargent, K.A., and Langer, W.H., 1989c, Studies of geology and hydrology in the Basin and Range province, southwestern United States, for isolation of high-level radioactive waste – evaluation of regions: U.S. Geological Survey Professional Paper 1370-H, 61 p. **(D1)**
- Bedinger, M.S., Sargent, K.A., and Reed, J.E., 1984, Geologic and hydrologic characterization of the Basin and Range province relative to the disposal of high-level radioactive waste – Part I, Introduction and guidelines: U.S. Geological Survey Circular 904-A, 16 p. **(D1)**
- Bedinger, M.S., Sargent, K.A., Langer, W.H., Sherman, K.B., Reed, J.E., and Brady, B.T., 1989, Studies of geology and hydrology in the Basin and Range province, southwestern United States, for isolation of high-level radioactive waste – Basis of characterization and evaluation: U.S. Geological Survey Professional Paper 1370-A, 41 p. **(D1)**
- Birkeland, P.W., 1984, Presentation of the Kirk Bryan Award to Leland H. Gile, John W. Hawley, and Robert B. Grossman, *in* Medals and Awards for 1983: Geological Society of America, v. 95, p. 1003-1004. **(A2)**
- Bowman, I., 1911, Well-drilling methods: USGS Water-Supply Paper 257, 139 p. **(D1)**
- Brand, D.C., 1937, The natural landscape of northwestern Chihuahua: University of New Mexico Press, 74 p. **(B2, C, C1, C2a, F3)**
- Bryan, K., 1923, Erosion and sedimentation in the Papago country, Arizona: U.S. Geological Survey, Bulletin 730-B, p. 19-90. **(C2a, A2)**
- Bryan, K., 1925, Date of channel trenching (arroyo cutting) in the arid Southwest: *Science*, n.s., v.62, p.338-344. **(C2a)**
- Bryan, K., 1936, Processes of formation of pediments at Granite Gap, New Mexico: *Zeitschrift für Geomorphologie*, Bd. 9, H. 4, p. 125-135. **(C2a)**
- Bryan, K., 1938, Geology and groundwater conditions of the Rio Grande depression in Colorado and New Mexico, *in* [U.S.] National Resources Committee, Regional Planning part VI – The Rio Grande joint investigations in the upper Rio Grande basin in Colorado, New Mexico, and Texas, 1936-1937: U.S. Government Printing Office, Washington, D.C., v. 1, part 2, p. 197-225. **(D1, G2, I3)**
- Bryan, K., 1940a, Erosion of the valleys of the southwest: *New Mexico Quarterly*, p. 227-232. **(C2a)**
- Bryan, K., 1940b, The retreat of slopes, *in* von Englen, O.D., ed., Symposium, Walther Penck's contribution to geomorphology: *Association of American Geographers Annals*, v. 30, p. 254-268. **(C2a)**
- Bryan, K., 1941, Correlation of the deposits of Sandia Cave, New Mexico, with the glacial chronology (appendix), *in* Hibben, F.C., *Evidences of Early Occupation in Sandia Cave, New Mexico, and Other Sites in the Sandia-Manzano Region*, Smithsonian Miscellaneous Collections 99, p. 45-64. **(B2, C1)**
- Bryan, K., and McCann, F.T., 1937, The Ceja del Rio Puerco: A border feature of the Basin and Range province in New Mexico, Part I. Stratigraphy and structure: *Journal of Geology*, v. 45, no. 8, p. 801-828. **(C2a)**
- Bryan, K., and McCann, F.T., 1938, The Ceja del Rio Puerco: A border feature of the Basin and Range province in New Mexico, Part II. Geomorphology: *Journal of Geology*, v. 46, p. 1-16. **(C2a)**
- Chapin, C.E., 1971, The Rio Grande rift, Part I: Modifications and additions: *Guidebook of the San Luis Basin, Colorado*, New Mexico Geological Society Guidebook 22, p. 191-201. **(C2a, I3)**
- Chapin, C.E., and Seager, W.R., 1975, Evolution of the Rio Grande rift in the Socorro and Las Cruces areas: *New Mexico Geological Society Guidebook 26*, p. 297-321. **(C2a, I3)**
- Cliett, T., 1969, Groundwater occurrence of the El Paso area and its related geology: *New Mexico Geological Society Guidebook 20*, p. 209-214. **(H1)**
- Cliett, T., and Hawley, J.W., 1996, General geology and groundwater occurrence of the El Paso area, *in* Ortega Klett, C.T., ed., *Proceedings of the 40th Annual New Mexico Water Conference*, New Mexico Water Resources Research Institute Report No. 297, p. 51-56. **(H1, H2)**
- Collins, E.W., and Raney, J.A., 2000, Geologic map of west Hueco Bolson, El Paso region, Texas: University of Texas at Austin, Bureau of Economic Geology, Miscellaneous Map No. 40, scale 1:100,000. **(C2b, F1)**
- Connell, S.D., Hawley, J.W., and Love, D.W., 2005, Late Cenozoic drainage development in the southeastern Basin and Range of New Mexico, southeasternmost Arizona and western Texas, *in* Lucas, S.G., Morgan, G., and Zeigler, K.E., eds., 2005, *New Mexico's Ice Ages*: New Mexico Museum of Natural History & Science Bulletin No. 28, p. 125-150. **(C2b, I1, I3)**
- Conover, C.S., 1954, Ground-water conditions in the Rincon and Mesilla Valleys and adjacent areas in New Mexico: U.S. Geological Survey Water-Supply Paper 1230, 200 p. **(G2)**

- Darton, N.H., 1899, Preliminary report on the geology and water resources of Nebraska west of the 103rd Meridian: U.S. Geological Survey 19th Annual Report, part 4, p. 719-785, plates 74-118. [Ogallala *lithostratigraphic unit* named] **(D1)**
- Darton, N.H., 1905, Preliminary report on the geology and underground water resources of the central Great Plains: U.S. Geological Survey Professional Paper 32, 433 p. [Ogallala *lithostratigraphic unit* defined] **(D1)**
- Darton, N.H., 1914, Underground water of Luna County, New Mexico: U.S. Geological Survey Water-Supply Paper 345, p. 25-40. **(D1, G1)**
- Darton, N.H., 1916a, Explosion craters: Science Monthly, v. 3, no. 4, p. 417-430 **(C2a, D1, G1)**
- Darton, N.H., 1916b, Geology and underground water of Luna County, New Mexico: U.S. Geological Survey Bulletin 618, 188 p. **(D1, G1)**
- Darton, N.H., 1917, Deming folio: U.S. Geological Survey Geologic Atlas of the United States, Folio 207, 15 p., 4 maps (scale 1:125,000), 9 Plates. **(C2a)**
- Darton, N.H., 1922, Geologic structure of parts of New Mexico: U.S. Geological Survey Bulletin 726-E, p. 173-275 **(C2a)**
- Darton, N.H., 1928a, Geologic map of New Mexico: U.S. Geological Survey, scale 1:500,000. **(C2a, D1)**
- Darton, N.H., 1928b, Red beds and associated formations within New Mexico with outline of the geology of the state: U.S. Geological Survey, Bulletin 794, 356 p. **(C2a, D1)**
- Darton, N.H., 1933, Guidebook of the western United States, Part F, The Southern Pacific Lines, New Orleans to Los Angeles. U.S. Geological Survey Bulletin 845. 304 p., 29 route maps. [West TX, and Southern NM and AZ: p. 120-160]. **(B3, C2a, F2, G1)**
- Darton, N.H., and others, 1916, Guidebook to the western United States, part C, the Santa Fe route: U.S. Geological Survey, Bulletin 613, p. 81-100. **(D1)**
- Davis, S.N., and Davis, A.G., 2005, Hydrogeology in the United States, 1780-1950: Tucson, Department of Hydrology and Water Resources, University of Arizona, Publication HWR No. 05-02. 140 p. **(A2)**
- Deming, D., 2002, Introduction to hydrogeology: New York, The McGraw-Hill Book Companies, Inc., 468 p. ISBN 0-07-232622-0 **(D1)**
- Follett, W.W., 1898, Rio Grande Waters - Equitable distribution of waters of the Rio Grande: Message to the President of the United States: 55th Congress, 2nd Session, U.S. Senate Document No. 229, Washington, D.C., US Government Printing Office, 289 p. **(B3, D1, E3)**
- Freethy, G.W., Pool, D.R., Anderson, T.W., and Tucci, P., 1986, Description and generalized description of aquifer materials in alluvial basin of Arizona and adjacent parts of California and New Mexico: U.S. Geological Survey Hydrologic Investigations Atlas HA-664, 4 sheets, scale 1:500,000. **(H1)**
- Frenzel, P.F., and Kaehler, C.A., 1990, Geohydrology and simulation of ground-water flow in the Mesilla Basin, Doña Ana County, New Mexico and El Paso County, Texas; *with a section on Water quality and geochemistry by S.K. Anderholm*: U.S. Geological Survey Open-File Report 88-305, 179 p. **(F2, H2, H3)**
- Frenzel, P.F., and Kaehler, C.A., 1992, Geohydrology and simulation of ground-water flow in the Mesilla Basin, Doña Ana County, New Mexico and El Paso County, Texas, *with a section on Water quality and geochemistry by S.K. Anderholm*: U.S. Geological Survey Professional Paper 1407-C, 105 p. **(F2, H2, H3)**
- Fryar, A.E., 2007, The future of hydrogeology, then and now: A look back at O.E. Meinzer's perspectives, 1934 to 1947: Ground Water, v. 45, no. 2, p. 246-249. **(A2, D1)**
- Gates, J.S., and Stanley, W.D., 1976, Hydrologic interpretation of geophysical data from the southeastern Hueco Bolson, El Paso and Hudspeth Counties, Texas: U.S. Geological Survey Open-File Report 76-650, 37 p. **(C4, F2, H1)**
- Gates, J.S., White, D.E., and Leggat, E.R., 1984, Preliminary study of aquifers of the lower Mesilla Valley in Texas and New Mexico by model simulation: U.S. Geological Survey Water Resources Investigations Report 84-4317, 21 p. **(H1, H3)**
- Gates, J.S., White, D.E., Stanley, W.D., and Ackermann, H.D., 1980, Availability of fresh and slightly saline ground water in the basins of westernmost Texas: Texas Department of Water Resources, Report 256, 108 p. [First published as USGS Open-File Report 78-663, 115 p.] **(E2a, H2)**
- Gile, L.H., 1961, A classification of ca horizons in soils of a desert region, Doña Ana County, New Mexico: Soil Science Society of America Proceedings, v. 25, no. 1, p. 52-61. **(C3)**
- Gile, L.H., and Grossman, R.B., 1968, Morphology of the argillic horizon in desert soils of southern New Mexico: Soil Science, v. 106, no. 1, p. 6-15. **(C3)**
- Gile, L.H., and Grossman, R.B., 1979, The Desert Project Soil Monograph: U.S. Department of Agriculture, National Technical Information Service, Document No. PB80-13534, Springfield, VA 22161, 984 p. **(C3)**

- Gile, L.H., and Hawley, J.W., 1966, Periodic sedimentation and soil formation on an alluvial-fan piedmont in southern New Mexico: *Soil Science Society of America Proceedings*, v. 30, p. 261-268. **(C2a, C3)**
- Gile, L.H., and Hawley, J.W., 1968, Age and comparative development of desert soils at the Gardner Spring radiocarbon site, New Mexico: *Soil Science Society of America Proceedings*, v. 32, no. 5, p. 709-719. **(C2a, C3)**
- Gile, L.H., and Hawley, J.W., 1972, The prediction of soil occurrence in certain desert regions of the southwestern United States: *Soil Science Society of America Proceedings*, v. 36, no. 1, p. 119-124. **(C2a, C3)**
- Gile, L.H., Hawley, J.W., and Grossman, R.B., 1981, Soils and geomorphology in the Basin Range area of southern New Mexico – Guidebook to the Desert Project: New Mexico Bureau of Mines and Mineral Resources, Memoir 39, 222 p. <https://geoinfo.nmt.edu/publications/monographs/memoirs/39/> **(C2a, C3)**
- Gile, L.H., Peterson, F.F., and Grossman, R.B., 1965, The K horizon--a master soil horizon of carbonate accumulation: *Soil Science*, v. 99, no. 2, p. 74-82. **(C3)**
- Gile, L.H., Peterson, F.F., and Grossman, R.B., 1966, Morphological and genetic sequences of carbonate accumulation in desert soils: *Soil Science*, v. 101, no. 5, p. 347-360. **(C3)**
- Gile, L.H., Grossman, R.B., Hawley, J.W., and Monger, H.C., 1996, Ancient soils of the Rincon surface, northern Doña County, in Gile, L.H., and Ahrens, R.J., eds., *Supplement to the Desert Project Soil Monograph, Volume II: Soils and landscapes astride the Rio Grande Valley near Las Cruces, New Mexico*: U.S. Department of Agriculture, Natural Resources Conservation Service, National Soil Survey Center, Lincoln, NE, Soil Survey Investigation Report No. 44, p. 1-110. **(C2b, C3)**
- Gile, L.H., Hawley, J.W., Grossman, R.B., Monger, H.C., Montoya, C.E., and Mack, G.H., 1995, Supplement to the Desert Project Guidebook, with emphasis on soil micromorphology: New Mexico Bureau of Mines and Mineral Resources, Bulletin 142, 96 p. **(C2b, C3)**
- Gross, J., and Icerman, L., 1983, Subsurface investigations for the area surrounding Tortugas Mountain, Doña Ana County, New Mexico: New Mexico Energy Research and Development Institute, Interim Report NMERDI 2-67-2238 (2), 70 p. **(C4, H1)**
- Hackett, O.M., 1964, The father of modern ground water hydrology: *Ground Water*, v. 2, no. 1, p. 2-5. **(A2, D1)**
- Hamilton, S.L., and Maddock, T. III, 1993, Application of a groundwater flow model to the Mesilla Basin, New Mexico and Texas: University of Arizona Department of Hydrology and Water Resources, no. 93–020, variously paginated. **(H3)**
- Hanson, R.T., McLean, J.S., and Miller, R.S., 1994, Hydrogeologic framework and preliminary simulation of ground-water flow in the Mimbres Basin, southwestern New Mexico: U.S. Geological Survey Water-Resources Investigations Report 94-4011, 118 p. **(F2, H1, H3)**
- Hanson, R.T., Ritchie, A.B., Boyce, S.E., Galanter, A.E., Ferguson, I.A., Flint, L.E., and Henson, W.R., 2018, Rio Grande transboundary integrated hydrologic model and water-availability analysis, New Mexico and Texas, United States, and Northern Chihuahua, Mexico: U.S Geological Survey Open-File Report 2018–1091, 185 p. **(F1, H3)**
- Harris, L.G., 2012, Whose water is it anyway? Anatomy of the water war between El Paso, Texas and New Mexico, in Ortega Klett, C.T., ed., *One hundred years of water wars in New Mexico*: Santa Fe, Sunstone Press, p. 227-253. ISBN 978-0-86524-902-5 **(A2, B3, E2, E3)**
- Hawley, J.W., 1969b, Notes on the geomorphology and late Cenozoic geology of northwestern Chihuahua: New Mexico Geological Society Guidebook 20, p. 131-142. **(C2a, F3)**
- Hawley, J.W., compiler, 1970, Cenozoic stratigraphy of the Rio Grande Valley area, Doña Ana County, New Mexico: El Paso Geological Society, Guidebook 4, 49 p. **(C2a, F2)**
- Hawley, J.W., 1972, Geologic-geomorphic mapping to serve soil resource development: *Soil Conservation Society of America, Proceedings of the 27th Annual Meeting*, p. 24-30. **(C2a, C3)**
- Hawley, J.W., 1975a, Quaternary history of Doña Ana County region, south-central New Mexico: New Mexico Geological Society Guidebook 26, p. 139-150. **(C2a, F1, I3)**
- Hawley, J.W., 1975b, The desert soil-geomorphology project: New Mexico Geological Society, Guidebook 26, p. 183-185. **(A2, C3)**
- Hawley, J.W., compiler, 1978, Guidebook to the Rio Grande rift in New Mexico and Colorado: New Mexico Bureau of Mines and Mineral Resources, Circular 163, 241 p. **(A2, C2a, C4)**
- Hawley, J.W., 1984, Hydrogeologic cross sections of the Mesilla Bolson, New Mexico and Texas: New Mexico Bureau of Mines and Minerals Resources, Open-File Report 190, 10 p. *Appendix in Peterson, D.M., Khaleel, R., and Hawley, J.W., 1984. (H1)*

- Hawley, J.W., 2005, Five million years of landscape evolution in New Mexico: An overview based on two centuries of geomorphic conceptual-model development, *in* Lucas, S.G., et al., eds., *New Mexico's Ice Ages: New Mexico Museum of Natural History & Science Bulletin No. 28*, p. 9-93. **(A1, A2, C2b, F1, H1, I3)**
- Hawley, J.W., 2014a, Biographical profile of Leland H. Gile, Jr., *in* Kues, B.S., Lewis, C.J., and Lueth, V.W., *A brief history of geological studies in New Mexico: New Mexico Geological Society, Special Publication 12*, p. 187-189 (*cited references 209-230*). **(A2)**
- Hawley, J.W., 2014b, Biographical profile of Oscar E. Meinzer, *in* Kues, B.S., Lewis, C.J., and Lueth, V.W., *A brief history of geological studies in New Mexico: New Mexico Geological Society, Special Publication 12*, p. 117-119 (*cited references 209-230*). **(A2)**
- Hawley, J.W., and Gile, L.H., 1966, Landscape evolution and soil genesis in the Rio Grande region, southern New Mexico: Friends of the Pleistocene, Rocky Mountain Section, Guidebook 11th Field Conference; La Cruces, New Mexico State University Agronomy Department, Special Publication, 74 p. **(C2a, C3)**
- Hawley, J.W., and Kennedy, J.F., 2004, Creation of a digital hydrogeologic framework model of the Mesilla Basin and southern Jornada del Muerto Basin: New Mexico Water Resources Research Institute Report No. 332, 105 p., with plates and appendix on CD ROM. <https://nmwrri.nmsu.edu/publications/technical-reports/tr-reports/tr-332.html> **(F1, H1)**
- Hawley, J.W., and Kernodle, J.M., 2000, Overview of the hydrogeology and geohydrology of the northern Rio Grande basin – Colorado, New Mexico, and Texas, *in* Ortega Klett, C.T., ed., *Proceedings of the 44th Annual New Mexico Water Conference: New Mexico Water Resources Research Institute Report No. 312*, p. 79-102. **(D1)**
- Hawley, J.W., and Kernodle, J.M., 2008, Early contributions to arid-zone hydrogeology in the eastern Basin and Range region: *Ground Water*, v. 46, no. 3, p. 510-516. **(A2, B3, D1)**
- Hawley, J.W., and Kottowski, F.E., 1969, Quaternary geology of the south-central New Mexico border region, *in* *Border Stratigraphy Symposium: New Mexico Bureau of Mines and Mineral Resources, Circular 104*, p. 89-115. **(C2a)**
- Hawley, J.W., and Lozinsky, R.P., 1992, Hydrogeologic framework of the Mesilla Basin in New Mexico and western Texas: New Mexico Bureau of Mines and Mineral Resources, Open-File Report 323, 55 p. **(H1)**
- Hawley, J.W., Kennedy, J.F., and Creel, B.J., 2001, The Mesilla Basin aquifer system of New Mexico, West Texas and Chihuahua – An overview of its hydrogeologic framework and related aspects of groundwater flow and chemistry, *in* Mace, R.E., Mullican, W.F. III, and Angle, E.S., eds., *Aquifers of West Texas: Texas Water Development Board Report 356*, p. 76-99. **(H1)**
- Hawley, J.W., Kennedy, J.F., Granados Olivas, A., and Ortiz, M.A., 2009, Hydrogeologic framework of the binational western Hueco Bolson-Paso del Norte area, Texas, New Mexico, and Chihuahua: Overview and progress report on digital model development: New Mexico Water Resources Research Institute Report No. 349, 45 p., 2 pls. <https://nmwrri.nmsu.edu/publications/technical-reports/tr-reports/tr-349.html> **(F1, H1)**
- Hawley, J.W., Kennedy, J.F., Ortiz, M.A., and Carrasco, S., 2005, Creation of a digital hydrogeologic framework model of the Rincon Valley and adjacent areas of Dona Ana, Sierra and Luna Counties NM: New Mexico Water Resources Research Institute, Addendum to Report No. 332 on CD ROM. <https://nmwrri.nmsu.edu/publications/technical-reports/tr-reports/tr-332.html> **(H1)**
- Hawley, J.W., Kottowski, F.E., Seager, W.R., King, W.E., Strain, W.S., and LeMone, D.V., 1969, The Santa Fe Group in the south-central New Mexico border region, *in* *Border Stratigraphy Symposium: New Mexico Bureau of Mines and Mineral Resources, Circular 104*, p. 52-76. **(C2a, F1)**
- Heath, R.C., 1988, Hydrogeologic setting of regions, *in* Back, W., Rosenshein, J.S., and Seaber, P.R., eds. *Hydrogeology – The Geology of North America: Geological Society of America, DNAG Volume. O-2*, p. 15-23. **(D1)**
- Hibbs, B.J., Creel, B.J., Boghici, R., Hayes, M., Ashworth, J., Hanson, A., Samani, Z., Kennedy, J.F., Hann, P., and Stevens, K., 1997, Transboundary Aquifers of the El Paso/Ciudad Juarez/Las Cruces Region: U.S. Environmental Protection Agency, Region 6; Technical Contract Report - Interagency Contracts X-996343-01-0 and X-996350-01-0, prepared by the Texas Water Development Board and the New Mexico Water Resources Research Institute, variously paged. *See TWDB, 1997, Appendix C–G.I.S. coverages, metadata descriptions, [and] groundwater data sets on CD-ROM, with Water Quality map insert on back-cover.* <https://nmwrri.nmsu.edu/publications/publications.html> **(F1, H1, H2)**
- Hill, R.T., 1891, Preliminary notes on the topography and geology of northern Mexico and southwest Texas, and New Mexico: *American Geologist*, v. 8, p. 133-141. **(B3, C2a, F1)**
- Hill, R.T., 1892b, Underground waters of the arid region: *Engineering Magazine*, v. 3, p. 653-660. **(G1)**

- Hill, R.T., 1900, Physical geography of the Texas region: U.S. Geological Survey Topographical Atlas Folio 3, 12 p. **(B3, C2a, G1)**
- Hoffer, J.M., 2001b, Geology of the West Potrillo Mountains, in Crumpler, L.S., and Lucas, S.G., eds., *Volcanology in New Mexico*: New Mexico Museum of Natural History and Science, Bulletin 18, p. 141-145. **(C2b)**
- Iceman, L., and Lohse, R.L., 1983, Geothermal low-temperature reservoir assessment in Dona Ana County, New Mexico: New Mexico Energy Research and Development Institute Report 2-69-2202, 188 p. **(C4, H2)**
- Jackson, D.B., 1976, Schlumberger soundings in the Las Cruces, New Mexico, area: U.S. Geological Survey Open-File Report 76-231, 170 p. **(C4)**
- Jackson, D.B., and Bisdorf, R.J., 1975, Direct-current soundings on the La Mesa Surface near Kilbourne and Hunt Holes, New Mexico: New Mexico Geological Society Guidebook 26, p. 273-275. **(C2a, C4)**
- Johnson, W.D., Jr., Hawley, J.W., Stone, W.J., Kottowski, F.E., Henry, C.D., and Price, J.G., 1989, Geology, in Bedinger, M.S., and others, *Studies of geology and hydrology in the Basin and Range Province, southwestern United States, for isolation of high-level radioactive waste; characterization of the Rio Grande region, New Mexico and Texas*: U.S. Geological Survey Professional Paper 1370-C, p. C7-C19. **(C2a)**
- Keller, G.R., and Cather, S.M., eds., 1994, Basins of the Rio Grande rift: Structure, stratigraphy and tectonic setting: Geological Society of America Special Paper 291, 304 p. **(C2b, C4)**
- Keller, G.R., Morgan, P., and Seager, W.R., 1990, Crustal structure, gravity anomalies and heat flow in the southern Rio Grande rift and their relationship to extensional tectonics: *Tectonophysics*, v. 174, p. 21-7. **(C2b, C4)**
- Kelley, V.C., 1952, Tectonics of the Rio Grande depression of central New Mexico: New Mexico Geological Society Guidebook 3, p. 92-105. **(C2a)**
- Kernodle, J.M., 1992a, Results of simulations by a preliminary numerical model of land subsidence in the El Paso, Texas, area: U.S. Geological Survey Water-Resources Investigations Report 92-4037, 35 p. **(D1, H3)**
- Kernodle, J.M., 1992b, Summary of U.S. Geological Survey ground-water-flow models of basin-fill aquifers in the southwestern alluvial basins region, Colorado, New Mexico, and Texas: U.S. Geological Survey Open-File Report 90-361, 81 p. **(D2, H3)**
- Khaleel, R., Palumbo, M.R., Peterson, D.D., and Hawley, J.W., 1983, Numerical Modeling of groundwater flow in the lower Rio Grande Basin, New Mexico: New Mexico Water Resources Research Institute, Project No. 135465 Completion Report, 188 p. **(H3)**
- King, W.E., and Hawley, J.W., 1975, Geology and ground-water resources of the Las Cruces area: New Mexico Geological Society Guidebook 26, p. 195-204. **(H1)**
- King, W.E., Hawley, J.W., Taylor, A.M., and Wilson, R.P., 1969, Hydrogeology of the Rio Grande Valley and adjacent intermontane areas of southern New Mexico: New Mexico Water Resources Research Institute Miscellaneous Report No. M6, 141 p. <https://nmwrri.nmsu.edu/publications/miscellaneous-reports/m-6/documents/m6.pdf> **(F2, H1, H2)**
- King, W.E., Hawley, J.W., Taylor, A.M., and Wilson, R.P., 1971, Geology and ground-water resources of central and western Doña Ana County, New Mexico: New Mexico Bureau of Mines and Mineral Resources, Hydrologic Report 1, 64 p. **(F2, H1, H2)**
- Knorr, D.B., and Cliett, T., 1985, Proposed groundwater recharge at El Paso, Texas, in Asano, T., ed., *Artificial recharge of groundwater*: Boston, Butterworth Publishers, p. 425-479. **(D2, E2b)**
- Knowles, D.B., and Kennedy, R.A., 1958, Ground-water resources of the Hueco Bolson northeast of El Paso, Texas: U.S. Geological Survey Water-Supply Paper 1426, 186 p. **(F2, G2)**
- Kottowski, F.E., 1958a, Geologic history of the Rio Grande near El Paso: West Texas Geological Society, Guidebook 1958 Field Trip, p. 46-54. *N.M. Bureau of Mines and Mineral Resources, Reprint Series No. 1. Reprint produced with permission of the West Texas Geological Society, Inc.* **(C2a, F1)**
- Kues, B.S., 2014a, Biographical profile of N.H. Darton, in Kues, B.S., Lewis, C.J., and Lueth, V.W., *A brief history of geological studies in New Mexico*: New Mexico Geological Society, Special Publication 12, p. 97-98 (cited references 209-230). **(A2)**
- Kues, B.S., 2014b, Biographical profile of Willis T. Lee, in Kues, B.S., Lewis, C.J., and Lueth, V.W., *A brief history of geological studies in New Mexico*: New Mexico Geological Society, Special Publication 12, p. 91-92 (cited references 209-230). **(A2)**
- Kues, B.S., Lewis, C.J., and Lueth, V.W., 2014, *A brief history of geological studies in New Mexico, with biographical Profiles of notable New Mexico geologists*: New Mexico Geological Society, Special Publication 12, 230 p. **(A2)**
- Lee, W.T., 1907a, Afton Craters of southern New Mexico: Geological Society of America Bulletin, v. 18, p. 211-220. **(C2a, F2, G1)**

- Lee, W.T., 1907b, Water resources of the Rio Grande Valley in New Mexico: U.S. Geological Survey Water-Supply Paper 188, 59 p. **(C2a, F2, G1)**
- Lee, W.T., 1922, The face of the earth as seen from the air: A study in the application of airplane photography to geography: American Geographical Society, Special Publication No. 1, 110 p. **(D1, E1)**
- Lee, W.T., 1924, A visit to Carlsbad Cavern: National Geographic Magazine, v. 45, no. 1, p. 1-40. **(D1)**
- Lee, W.T., 1925a, Erosion by solution and fill: U.S. Geological Survey Bulletin 760-D, p. 107-121. **(D1)**
- Lee, W.T., 1925b, New discoveries in Carlsbad Cavern: National Geographic Magazine, v. 48, no. 3, p. 301-319. **(D1)**
- Leggat, E.R., Lowry, M.E., and Hood, J.W., 1962, Ground-water resources of the lower Mesilla Valley, Texas and New Mexico: Texas Water Commission, Bulletin 6203, 191 p. **(F2, G2, H1, H2)**
- Leopold, L.B., 2004, Geomorphology: A sliver off the corpus of science: Annual Review of Earth and Planetary Science, v. 32, p. 1-12. **(A2)**
- Mace, R.E., Mullican, W.F. III, and Angle, E.S., eds., 2001, Aquifers of West Texas: Texas Water Development Board Report 356, 272 p. **(D1, F2)**
- Mack, G.H., Kottlowski, F.E., and Seager, W.R., 1998, The stratigraphy of south-central New Mexico: New Mexico Geological Society Guidebook 49, p. 135-154. **(C2b, I3)**
- Mackichan, L., 1980, Clyde Wilson Dies at Age 48: The Divining Rod, New Mexico Water Resources Research Institute, no. 80-3 (Fall 1980), p. 1, 3. <https://nmwrri.nmsu.edu/publications/enews-divining-rod-issues/enews-divining-rod-issues-documents/Vol.%2080%20No.%203%20-%20Fall%201980.pdf> **(A2)**
- Mattick, R.E., 1967, A seismic and gravity profile across the Hueco bolson, Texas: U.S. Geological Survey Professional Paper 575-D, p. D85-D91. **(C2a, C4)**
- Maxey, G.B., 1979, The Meinzer era of U.S. hydrogeology, 1910-1940, in Back, W., and Stephenson, D.A., eds., Contemporary hydrology – The George Burke Maxey Memorial Volume: Journal of Hydrology, v. 43, no. 1/6, p. 1-6. **(A2)**
- Meinzer, O.E., 1911, Geology and water resources of Estancia Valley, New Mexico: U.S. Geological Survey Water-Supply Paper 275, 89 p. **(B2, C1, C2a, D1, G1, I1, I2)**
- Meinzer, O.E., 1922, Map of the Pleistocene lakes of the Basin and Range province and its significance: Geological Society of America Bulletin, v. 33, issue 3, p. 541-552. **(B2, C1, D1, I1)**
- Meinzer, O.E., 1923b, Outline of ground water hydrology, with definitions: U.S. Geological Survey Water-Supply Paper 494, 71 p. **(A1, D1)**
- Meinzer, O.E., and Hare, R.F., 1915, Geology and water resources of the Tularosa Basin, New Mexico: U.S. Geological Survey Water-Supply Paper 343, 317 p. **(B2, C1, C2a, D1, G1, I2)**
- Meinzer, O.E., and Kelton, F.C., 1912, Geology and water resources of Sulphur Springs Valley, Arizona: U.S. Geological Survey Water-Supply Paper 320, 231 p. **(B2, C1, D1, G1, I1)**
- Meyer, G., Davis, G., and La Moreaux, P.E., 1988, Historical perspective - Chapter 1; in Back, W., Rosenshein, J.S., and Seaber, P.R., eds., Geology of North America – Hydrogeology: Geological Society of America, Decade of North American Geology, v. 0-2, p. 1-8. **(A2, D1)**
- Monger, H.C., Gile, L.H., Hawley, J.W., and Grossman, R.B., 2009, The Desert Project – An analysis of aridland soil-geomorphic processes: New Mexico State University Agricultural Experiment Station Bulletin 798, 76 p. **(A2, C2b, C3)**
- Myers, R.G., and Orr, B.R., 1986, Geohydrology of the aquifer in the Santa Fe Group, northern West Mesa of the Mesilla Basin near Las Cruces, New Mexico: U.S. Geological Survey Water-Resources Investigations Report 84-4190, 37 p. **(C2a, H1)**
- NACOSN (North American Commission on Stratigraphic Nomenclature), 2005, North American Stratigraphic Code: The American Association of Petroleum Geologists Bulletin, v. 89, p. 1547-1591. **(B1)**
- Newell, F.H., 1893, Water supply for irrigation: U.S. Geological Survey, 13th Annual Report 13, Part 3a, p. 7-99. **(B3, D1)**
- Nickerson, E.L., 1989, Aquifer tests in the flood-plain alluvium and Santa Fe Group at the Rio Grande near Canutillo, El Paso County, Texas: U.S. Geological Survey Water-Resources Investigations Report 89-4011, 29 p. **(H3)**
- Nickerson, E.L., 1995, Selected geohydrologic data for the Mesilla ground-water basin, 1987 through 1992 water years, Doña Ana County, New Mexico and El Paso County, Texas: U.S. Geological Survey Open-File Report 95-111, 123 p. **(H1)**
- Nickerson, E.L., 2006, Description of piezometers and ground-water-quality characteristics at three new sites in the Lower Mesilla Valley, Texas, and New Mexico: U.S. Geological Survey Scientific Investigations Report 2005-5248, 27 p. **(H1, H2)**

- Nickerson, E.L., and Myers, R.G., 1993, Geohydrology of the Mesilla ground-water basin, Doña Ana County, New Mexico, and El Paso County, Texas: U.S. Geological Survey Water-Resources Investigations Report 92-4156, 89 p. **(H1, H2)**
- Orr, B.R., and Risser, D.W., 1992, Geohydrology and potential effects of development of freshwater resources in the northern part of the Hueco Bolson, Doña Ana and Otero Counties, New Mexico, and El Paso County, Texas: U.S. Geological Survey Water-Resources Investigations Report 91-4082, 92 p. **(H1, H2)**
- Paige, S., and Darton, N.H., 1916, Descriptions of the Silver City Quadrangle, New Mexico: United States Geological Survey, Geologic Atlas Folio 199, 10 p. **(C2a, G1)**
- Peterson, D.M., Khaleel, R., and Hawley, J.W., 1984, Quasi three-dimensional modeling of groundwater flow in the Mesilla Bolson, New Mexico: New Mexico Water Resources Research Institute Report No. 178, 185 p. <https://nmwrrr.nmsu.edu/publications/technical-reports/tr-reports/tr-178.html> **(H1, H3)**
- Phillips, F.M., 2014, Biographical profile of Kirk Bryan, *in* Kues, B.S., Lewis, C.J., and Lueth, V.W., A brief history of geological studies in New Mexico: New Mexico Geological Society, Special Publication 12, p. 142-143 (*cited references 209-230*). **(A2)**
- Rabbitt, M.C., 1989, The United States Geological Survey: 1879 to 1989: U.S. Geological Survey, Circular 1050, 52 p. **(A2)**
- Reeder, H.O., 1957, Ground water in Animas Valley, Hidalgo County, New Mexico: New Mexico State Engineer Office, Technical Report No. 11, 101 p. **(G2, I2)**
- Reynolds, C.B. & Associates, 1986, Shallow seismic reflection survey, Canutillo area, Texas and New Mexico: Unpublished consultant's report for U.S. Geological Society by Charles B. Reynolds & Associates, 4409 San Andres Ave., NE, Albuquerque, NM 87110, 8 p., 2 figs., 12 enclosed seismic profiles. **(C4)**
- Reynolds, C.B. & Associates, 1987, Shallow seismic reflection line CA-1, Doña Ana County, New Mexico and El Paso County, Texas: Unpublished consultant's report for U.S. Geological Society by Charles B. Reynolds & Associates, 4409 San Andres Ave., NE, Albuquerque, NM 87110, 4 p., 1 fig., 2 enclosed seismic profiles. **(C4)**
- Richardson, G.B., 1909, Description of the El Paso quadrangle, Texas: U.S. Geological Survey Geological Atlas, El Paso folio, ser. no. 116, 11 p. **(G1)**
- Richardson, G.L., Gebbard, T.G., Jr., and Brutsaert, W.F., 1972, Water-table investigation in the Mesilla Valley: Las Cruces, New Mexico State University, Engineering Experiment Station Technical Report 76, 206 p. **(H3)**
- Riecker, R.E., ed., 1979, Rio Grande rift: Tectonics and magmatism: Washington, D.C., American Geophysical Union, 438 p. **(A2, C2a, C4)**
- Ruhe, R.V., 1962, Age of the Rio Grande Valley in southern New Mexico, *Journal of Geology*, v. 70, p. 151-167. **(C2a)**
- Ruhe, R.V., 1964, Landscape morphology and alluvial deposits in southern New Mexico: *Annals of the Association of American Geographers*, v. 54, p. 147-159. **(C2a, C3)**
- Ruhe, R.V., 1967, Geomorphic surfaces and surficial deposits in southern New Mexico: New Mexico Bureau of Mines and Mineral Resources, Memoir 18, 65 p. **(C2a, C3)**
- Sargent, K.A., Hawley, J.W., Henry, C.D., Stone, W.J., and Kottowski, F.E., 1989, Quaternary tectonism, *in* studies of geology and hydrology in the Basin and Range Province, southwestern United States, for isolation of high-level radioactive waste; characterization of the Rio Grande Region, New Mexico and Texas: U.S. Geological Survey Professional Paper 1370-C, p. C23-C26. **(C2a)**
- Sayre, A.N., and Livingston, P., 1945, Ground-water resources of the El Paso area, Texas: U.S. Geological Survey Water-Supply Paper 919, 190 p. **(D1, F2, G2)**
- Sayre, A.N., and Stephenson, E.L., 1937, The use of resistivity methods in the location of salt-water bodies in the El Paso, Texas, area: *American Geophysical Union Transactions*, 18th Annual Meeting, p. 393-398. **(C4, G2, H2)**
- Schwennesen, A.T., 1918, Ground water in the Animas, Playas, Hachita, and San Luis Basins, New Mexico: U.S. Geological Survey Water-Supply Paper 422, 152 p. **(D1, F2, G1, I2)**
- Seager, W.R., 1981, Geology of the Organ Mountains and southern San Andres Mountains, New Mexico: New Mexico Bureau of Mines and Mineral Resources, Memoir 36, 97 p. **(C2a)**
- Seager, W.R., 1995, Geology of southwest quarter of Las Cruces and northwest El Paso 1° x 2° sheets: New Mexico Bureau of Mines and Mineral Resources, Geologic Map 60, scale 1:125,000. **(C2b, C4)**
- Seager, W.R., and Hawley, J.W., 1973, Geology of Rincon quadrangle, New Mexico: New Mexico Bureau of Mines and Mineral Resources, Bulletin 101, 56 p., map scale 1:24,000. **(C2a)**
- Seager, W.R., and Mack, G.H., 1994, Geology of the East Potrillo Mountains and vicinity, Doña Ana County, New Mexico: New Mexico Bureau of Mines and Mineral Resources, Bulletin 113, 27 p. **(C2b)**

- Seager, W.R., and Morgan, P., 1979, Rio Grande rift in southern New Mexico, west Texas and northern Chihuahua, *in* Riecker, R.E., ed., Rio Grande rift, tectonics and magmatism: Washington D.C., American Geophysical Union, p. 87-106. **(C2a, C4)**
- Seager, W.R., Clemons, R.E., and Hawley, J.W., 1975, Geology of Sierra Alta Quadrangle, Doña Ana County, New Mexico, New Mexico Bureau of Mines and Mineral Resources, Bulletin 102, 56 p., map scale 1:24,000. **(C2a)**
- Seager, W.R., Hawley, J.W., and Clemons, R.E., 1971, Geology of San Diego Mountain area, Doña Ana County, New Mexico: New Mexico Bureau of Mines and Mineral Resources, Bulletin 97, 38 p., map scale 1:24,000. **(C2)**
- Seager, W.R., Hawley, J.W., Kottowski, F.E., and Kelley, S.A., 1987, Geologic map of east half of Las Cruces and northeast El Paso 1° x 2° sheets, New Mexico: New Mexico Bureau of Mines and Mineral Resources, Geologic Map, GM-57, scale 1:125,000, 3 sheets. **(C2a)**
- Sharp, R.P., 1993, Recollections of Kirk Bryan: a biographical sketch: *Geomorphology*, v. 6, p. 189-205. **(A2)**
- Siebenthal, C.E., 1910, Geology and water resources of the San Luis Valley, Colorado: U.S. Geological Survey, Water-Supply Paper 240, 128 p. **(D1, I1, I3)**
- Slichter, C.S., 1899, Theoretical investigation of the motion of ground waters: U.S. Geological Survey Annual Report 19, part 2, p. 295-384. **(D2)**
- Slichter, C.S., 1902, The motions of underground water: U.S. Geological Survey Water-Supply and Irrigation Paper 67, 106 p. **(D2)**
- Slichter, C.S., 1905a, Field measurements of the rate of movement of underground waters: U.S. Geological Survey Water-Supply and Irrigation Paper 140, 122 p. **(D2, G1)**
- Slichter, C.S., 1905b, Observations on ground waters of the Rio Grande valley: U.S. Geological Survey Water-Supply and Irrigation Paper 141, 83 p. **(G1)**
- Spiegel, Z.E., 1962, Hydraulics of certain stream-connected aquifer systems: New Mexico State Engineer Special Report, 105 p. **(D2, G2)**
- Spiegel, Z.E., and Baldwin, B., 1963, Geology and water resources of the Santa Fe area, New Mexico: U.S. Geological Survey, Water-Supply Paper 1525, 258 p. **(D1, G2)**
- Stone, G.H., 1901, Notes on the extinct glaciers of Arizona and New Mexico: *Science*, v. 14, p. 798. **(B2, C1)**
- Strain, W.S., 1966, Blancan mammalian fauna and Pleistocene formations, Hudspeth County, Texas: Texas Memorial Museum, Bulletin 10, 55 p. **(B1, C2a)**
- Strain, W.S., 1971, Late Cenozoic bolson integration in the Chihuahua tectonic belt, *in* The geologic framework of the Chihuahua tectonic belt: West Texas Geological Society, Publication No. 71-59, p. 167-173. **(C2a)**
- Sweetkind, D.S., 2017, Three-dimensional hydrogeologic framework model of the Rio Grande transboundary region of New Mexico and Texas, USA, and northern Chihuahua, Mexico: U.S. Geological Survey Scientific Investigations Report 2017-5060, 49 p. **(H1)**
- Sweetkind, D.S., Hanson, R.T., Ritchie, A.B., and Hawley, J.W., 2017, Data release of Three-Dimensional Hydrogeologic Framework Model of the Rio Grande Transboundary Region of New Mexico and Texas, USA and Northern Chihuahua, Mexico: U.S. Geological Survey data release. **(H1)**
- Taylor, A.M., 1967, Geohydrologic investigations in the Mesilla Valley, New Mexico: New Mexico State University, master's thesis, 130 p. **(H1, H3)**
- Teeple, A.P., 2017a, Geophysics- and geochemistry-based assessment of the geochemical characteristics and groundwater-flow system of the U.S. part of the Mesilla Basin/Conejos-Médanos aquifer system in Doña Ana County, New Mexico, and El Paso County, Texas, 2010–12: U.S. Geological Survey Scientific Investigations Report 2017-5028, 183 p. **(C4, H2)**
- Theis, C.V., 1935a, The relation between lowering the piezometric surface and the rate and duration of discharge of well using ground water storage: *American Geophysical Union Transactions*, 16th Annual Meeting, part 2, p. 519-524. **(D1, D2)**
- Theis, C.V., 1935b, The source of water derived from wells: Essential factors controlling the response of an aquifer to development: *Civil Engineer*, v. 10, p. 277-280. **(D1, D2)**
- Theis, C.V., 1938, Ground water in the middle Rio Grande valley in [U.S.] Natural Resources Committee, Regional Planning part VI---The Rio Grande joint investigations in the upper Rio Grande basin in Colorado, New Mexico, and Texas, 1936-1937: Washington, D.C., U.S. Government Printing Office, v. 1, part 2, p. 268-291. **(D1, D2)**
- Tolman, C.F., 1909, Erosion and deposition in southern Arizona bolson region: *Journal of Geology*, v. VII, no. II, p. 136-163. **(C2a, D1)**
- Tolman, C.F., 1937, *Ground Water*: New York, McGraw-Hill Book Co., Inc. 593 p. **(A2, C, D1)**

- Trauger, F.D., 1972, Water Resources and General Geology of Grant County, New Mexico; Hydrologic Report 2; New Mexico Bureau of Mines and Mineral Resources, Socorro, 1972, 211 p. **(D1, H1, H2)**
- Trauger, F.D., and Doty, G.C., 1965, Ground water – Its occurrence and relation to the economy and geology of southwestern New Mexico: New Mexico Geological Society Guidebook 16, p. 215-227. **(E2, F2, G2)**
- U.S. National Resources Committee, 1938, The Rio Grande Joint Investigation in the Upper Rio Grande Basin in Colorado, New Mexico, and Texas: Washington, D.C., U.S. Government Printing Office. 566 p. **(B3, D1, G1)**
- Vanden Heuvel, R.C., 1966, The occurrence of sepiolite and attapulgite in the calcareous zone of a soil near Las Cruces, New Mexico, *in* Clays and clay minerals: Proceedings of the 13th National Conference on clays and clay minerals, New York, Pergamon Press, p. 193-207. **(C3)**
- Waring, G.A., and Meinzer, O.E., 1947, Bibliography and index of publications relating to ground water prepared by the Geological Survey and cooperating agencies: U.S. Geological Survey Water-Supply Paper 992, 412 p. **(A1)**
- White, W.N., 1931, Preliminary report on ground-water supply of Mimbres Valley, New Mexico: U.S. Geological Survey Water-Supply Paper 637, part b, p. 69-90. **(G1)**
- Wilkins, D.W., 1986, Geohydrology of the Southwest Alluvial Basins, Regional Aquifer-systems analysis in parts of Colorado, New Mexico, and Texas: U.S. Geological Survey Water Resources Investigations Report 84-4224, 61 p. **(D1, F2)**
- Wilkins, D.W., 1998, Summary of the southwest alluvial basins regional aquifer-system analysis in parts of Colorado, New Mexico, and Texas: U.S. Geological Survey Professional Paper 1407-A, 49 p. **(D1, F2)**
- Wilson, C.A., and White, R.R., 1984, Geohydrology of the central Mesilla Valley, Doña Ana County, New Mexico: U.S. Geological Survey Water Resources Investigations Report 82-444, 144 p. **(H1, H2)**
- Wilson, C.A., White, R.R., Orr, B.R., and Roybal, R.G., 1981, Water resources of the Rincon and Mesilla Valleys and adjacent areas, New Mexico: New Mexico State Engineer Technical Report 43, 514 p. **(H1, H2)**
- Woodward, L.A., Callender, J.F., Seager, W.R., Chapin, C.E., Gries, J.C., Schaffer, W.L., and Zilinski, R.E., 1978, Tectonic map of the Rio Grande rift region in New Mexico, Chihuahua, and Texas, *in* Hawley, J.W., compiler, Guidebook to the Rio Grande rift in New Mexico and Colorado: New Mexico Bureau of Mines and Mineral Resources Circular 163, Sheet 2, approx. scale 1:1,000,000. **(C2a)**
- Zohdy, A.A.R., Bisdorf, R.J., and Gates, J.S., 1976, Schlumberger soundings in the lower Mesilla Valley of the Rio Grande, Texas and New Mexico: U.S. Geological Survey Open-File Report 76-324, 77 p. *See Al-Garni 1996.* **(C4)**

C7.2. Chronology of Notable Contributions to the Understanding of the Binational Mesilla Basin Region's Hydrogeology and Geohydrology (1890-2006)

C7.2.1. 1890 to 1945

- Tarr, R.S., 1890, Drainage systems of New Mexico: American Geologist, v. 5, p. 261-270. **(C, G1)**
- Hill, R.T., 1891, Preliminary notes on the topography and geology of northern Mexico and southwest Texas, and New Mexico: American Geologist, v. 8, p. 133-141. **(B3, C2a, F1)**
- Hill, R.T., 1892a, Notes on the Texas-New Mexican region: Geological Society of America Bulletin, v. 3, p. 85-100. **(B3, C2a)**
- Hill, R.T., 1892b, Underground waters of the arid region: Engineering Magazine, v. 3, p. 653-660. **(G1)**
- Hill, R.T., 1896, Descriptive topographic terms of Spanish America: National Geographic, v. 7, p. 291-302. **(A1, B3, C2a, F1)**
- Barker, F.C., 1898, Irrigation in Mesilla Valley, New Mexico: U.S. Geological Survey Water-Supply and Irrigation Paper 10, 51 p. **(G1)**
- Follett, W.W., 1898, Rio Grande Waters - Equitable distribution of waters of the Rio Grande: Message to the President of the United States: 55th Congress, 2nd Session, U.S. Senate Document No. 229, Washington, D.C., US Government Printing Office, 289 p. **(B3, D1, E3)**
- Hill, R.T., 1900, Physical geography of the Texas region: U.S. Geological Survey Topographical Atlas Folio 3, 12 p. **(B3, C2a, G1)**
- Slichter, C.S., 1899, Theoretical investigation of the motion of ground waters: U.S. Geological Survey Annual Report 19, part 2, p. 295-384. **(D2)**
- Slichter, C.S., 1902, The motions of underground water: U.S. Geological Survey Water-Supply and Irrigation Paper 67, 106 p. **(D2)**

- Slichter, C.S., 1905a, Field measurements of the rate of movement of underground waters: U.S. Geological Survey Water-Supply and Irrigation Paper 140, 122 p. **(D2, G1)**
- Slichter, C.S., 1905b, Observations on ground waters of the Rio Grande valley: U.S. Geological Survey Water-Supply and Irrigation Paper 141, 83 p. **(G1)**
- Lee, W.T., 1907a, Afton Craters of southern New Mexico: Geological Society of America Bulletin, v. 18, p. 211-220. **(C2a, F2, G1)**
- Lee, W.T., 1907b, Water resources of the Rio Grande Valley in New Mexico: U.S. Geological Survey Water-Supply Paper 188, 59 p. **(C2a, F2, G1)**
- Richardson, G.B., 1909, Description of the El Paso quadrangle, Texas: U.S. Geological Survey Geological Atlas, El Paso folio, ser. no. 116, 11 p. **(G1)**
- Bowman, I., 1911, Well-drilling methods: USGS Water-Supply Paper 257, 139 p. **(D1)**
- Meinzer, O.E., and Kelton, F.C., 1912, Geology and water resources of Sulphur Springs Valley, Arizona: U.S. Geological Survey Water-Supply Paper 320, 231 p. **(B2, C1, D1, G1, I1)**
- Nelson, J.W., and Holmes, L.C., 1914, Soil Survey of the Mesilla Valley, New Mexico-Texas: U.S. Department of Agriculture, Bureau of Soils, Washington, D.C., U.S. Government Printing Office, 39 p. map scale 1:63,360. *Shallow depth to water table shown in large no. soil-test borings.* **(C3, G1)**
- Darton, N.H., 1914, Underground water of Luna County, New Mexico: U.S. Geological Survey Water-Supply Paper 345, p. 25-40. **(D1, G1)**
- Follansbee, R., Porter, E.A., Follett, W.W., and Gray, G.A., 1915, Water resources of the Rio Grande basin, 1888-1913: U.S. Geological Survey Water-Supply Paper 358, p. 679-694. **(G1)**
- Meinzer, O.E., and Hare, R.F., 1915, Geology and water resources of the Tularosa Basin, New Mexico: U.S. Geological Survey Water-Supply Paper 343, 317 p. **(B2, C1, C2a, D1, G1, I2)**
- Darton, N.H., 1916a, Explosion craters: Science Monthly, v. 3, no. 4, p. 417-430 **(C2a, D1, G1)**
- Darton, N.H., 1916b, Geology and underground water of Luna County, New Mexico: U.S. Geological Survey Bulletin 618, 188 p. **(D1, G1)**
- Paige, S., and Darton, N.H., 1916, Descriptions of the Silver City Quadrangle, New Mexico: United States Geological Survey, Geologic Atlas Folio 199, 10 p. **(C2a, G1)**
- Darton, N.H., 1917, Deming folio: U.S. Geological Survey Geologic Atlas of the United States, Folio 207, 15 p., 4 maps (scale 1:125,000), 9 Plates. **(C2a)**
- Schwennesen, A.T., 1918, Ground water in the Animas, Playas, Hachita, and San Luis Basins, New Mexico: U.S. Geological Survey Water-Supply Paper 422, 152 p. **(D1, F2, G1, I2)**
- White, W.N., 1931, Preliminary report on ground-water supply of Mimbres Valley, New Mexico: U.S. Geological Survey Water-Supply Paper 637, part b, p. 69-90. **(G1)**
- Darton, N.H., 1933, Guidebook of the western United States, Part F, The Southern Pacific Lines, New Orleans to Los Angeles. U.S. Geological Survey Bulletin 845. 304 p., 29 route maps. [West TX, and Southern NM and AZ: p. 120-160]. **(B3, C2a, F2, G1)**
- Dunham, K.C., 1935, The geology of the Organ Mountains, with an account of the geology and mineral resources of Dona Ana County, New Mexico: New Mexico Bureau of Mines and Mineral Resources, Bulletin 11, 272 p. **(C2a, G1)**
- Ordóñez, E., 1936, Physiographic provinces of Mexico: American Association of Petroleum Geologists Bulletin, v. 20, no. 10, p. 1277-1307. **(C, F3)**
- Sayre, A.N., and Stephenson, E.L., 1937, The use of resistivity methods in the location of salt-water bodies in the El Paso, Texas, area: American Geophysical Union Transactions, 18th Annual Meeting, p. 393-398. **(C4, G2, H2)**
- Bryan, K., 1938, Geology and groundwater conditions of the Rio Grande depression in Colorado and New Mexico, in [U.S.] National Resources Committee, Regional Planning part VI – The Rio Grande joint investigations in the upper Rio Grande basin in Colorado, New Mexico, and Texas, 1936-1937: U.S. Government Printing Office, Washington, D.C., v. 1, part 2, p. 197-225. **(D1, G2, I3)**
- Sayre, A.N., and Livingston, P., 1945, Ground-water resources of the El Paso area, Texas: U.S. Geological Survey Water-Supply Paper 919, 190 p. **(D1, F2, G2)**

C7.2.2 1946 to 1980

- Waring, G.A., and Meinzer, O.E., 1947, Bibliography and index of publications relating to ground water prepared by the Geological Survey and cooperating agencies: U.S. Geological Survey Water-Supply Paper 992, 412 p. **(A1)**

- Scalapino, R.A., 1949, Ground-Water Resources of the El Paso Area, Texas: Texas Board of Water Engineers, Progress Report No. 6, 22 p. **(D1, G2)**
- Conover, C.S., 1954, Ground-water conditions in the Rincon and Mesilla Valleys and adjacent areas in New Mexico: U.S. Geological Survey Water-Supply Paper 1230, 200 p. **(G2)**
- Jackson, D.B., 1976, Schlumberger soundings in the Las Cruces, New Mexico, area: U.S. Geological Survey Open-File Report 76-231, 170 p. **(C4)**
- Jackson, D.B., and Bisdorf, R.J., 1975, Direct-current soundings on the La Mesa Surface near Kilbourne and Hunt Holes, New Mexico: New Mexico Geological Society Guidebook 26, p. 273-275. **(C2a, C4)**
- Knowles, D.B., and Kennedy, R.A., 1958, Ground-water resources of the Hueco Bolson northeast of El Paso, Texas: U.S. Geological Survey Water-Supply Paper 1426, 186 p. **(F2, G2)**
- Kottlowski, F.E., 1958a, Geologic history of the Rio Grande near El Paso: West Texas Geological Society, Guidebook 1958 Field Trip, p. 46-54. *N.M. Bureau of Mines and Mineral Resources, Reprint Series No. 1. Reprint produced with permission of the West Texas Geological Society, Inc.* **(C2a, F1)**
- Kottlowski, F.E., 1958b, Lake Otero-second phase formation of New Mexico's gypsum dunes: Geological Society of America Bulletin, v. 33, p. 541-552. **(C2a, I2)**
- Spiegel, Z.E., 1962, Hydraulics of certain stream-connected aquifer systems: New Mexico State Engineer Special Report, 105 p. **(D2, G2)**
- Leggat, E.R., Lowry, M.E. and Hood, J.W., 1962, Ground-water resources of the lower Mesilla Valley, Texas and New Mexico: Texas Water Commission, Bulletin 6203, 191 p. **(F2, G2, H1, H2)**
- Leggat, E.R., Lowry, M.E., and Hood, J.W., 1963, Ground-water resources of the lower Mesilla Valley, Texas and New Mexico: U.S. Geological Survey, Water-Supply Paper 1669AA, 49 p. **(F2, G2, H1, H2)**
- Albritton, C.C., Jr., and Smith, J.F., Jr., 1965, Geology of the Sierra Blanca area, Hudspeth County, Texas: U.S. Geological Survey Professional Paper 594-J, 131 p. **(C2a, D1, F1)**
- Mattick, R.E., 1967, A seismic and gravity profile across the Hueco bolson, Texas: U.S. Geological Survey Professional Paper 575-D, p. D85-D91. **(C2a, C4)**
- Cliett, T., 1969, Groundwater occurrence of the El Paso area and its related geology: New Mexico Geological Society Guidebook 20, p. 209-214. **(H1)**
- Córdoba, D.A., Wengerd, S.A., and Shomaker, J.W., eds., 1969, Guidebook of the Border Region: New Mexico Geological Society Guidebook 20, 218 p. *See Road Log Section, p. 1-38.* **(C2a)**
- Hawley, J.W., 1969a, Geology and its relation to the hydrologic system, in King, W.E., Hawley, J.W., Taylor, A.M., and Wilson, R.P., Hydrogeology of the Rio Grande Valley and adjacent intermontane areas of southern New Mexico: New Mexico Water Resources Research Institute Report No. 6, p. 19-47; *cf. King, W.E., and others, Geology and ground-water resources of central and western Doña Ana County, New Mexico: New Mexico Bureau of Mines and Mineral Resources, Hydrologic Report 1, p. 9-24, 61-62.* **(C2a, H1)**
- Hawley, J.W., 1969b, Notes on the geomorphology and late Cenozoic geology of northwestern Chihuahua: New Mexico Geological Society Guidebook 20, p. 131-142. **(C2a, F3)**
- Hawley, J.W., Kottlowski, F.E., Seager, W.R., King, W.E., Strain, W.S., and LeMone, D.V., 1969, The Santa Fe Group in the south-central New Mexico border region, in Border Stratigraphy Symposium: New Mexico Bureau of Mines and Mineral Resources, Circular 104, p. 52-76. **(C2a, F1)**
- Zohdy, A.A.R., 1969, The use of Schlumberger and equatorial soundings in groundwater investigations near El Paso, Texas: Geophysics, v. 34, p. 713-728. **(C4)**
- Zohdy, A.A.R., Jackson, D.B., Mattick, R.E., and Peterson, D.L., 1969, Geophysical survey for ground water at White Sands Missile Range, New Mexico: U.S. Geological Survey Open-File Report 69-326, 144 p. **(C4)**
- King, W.E., Hawley, J.W., Taylor, A.M., and Wilson, R.P., 1971, Geology and ground-water resources of central and western Doña Ana County, New Mexico: New Mexico Bureau of Mines and Mineral Resources, Hydrologic Report 1, 64 p. **(F2, H1, H2)**
- Richardson, G.L., Gebbard, T.G., Jr., and Brutsaert, W.F., 1972, Water-table investigation in the Mesilla Valley: Las Cruces, New Mexico State University, Engineering Experiment Station Technical Report 76, 206 p. **(H3)**
- Gates, J.S., and Stanley, W.D., 1976, Hydrologic interpretation of geophysical data from the southeastern Hueco Bolson, El Paso and Hudspeth Counties, Texas: U.S. Geological Survey Open-File Report 76-650, 37 p. **(C4, F2, H1)**
- Hawley, J.W., Bachman, G.O., and Manley, K., 1976, Quaternary stratigraphy in the Basin and Range and Great Plains provinces, New Mexico and western Texas, in Mahaney, W.C., ed., Quaternary stratigraphy of North America: Stroudsburg, PA, Dowden, Hutchinson, and Ross, Inc., p. 235-274. **(A1, A2, D1, I1, I3)**
- Jackson, D.B., 1976, Schlumberger soundings in the Las Cruces, New Mexico, area: U.S. Geological Survey Open-File Report 76-231, 170 p. **(C4)**

- Zohdy, A.A.R., Bisdorf, R.J., and Gates, J.S., 1976, Schlumberger soundings in the lower Mesilla Valley of the Rio Grande, Texas and New Mexico: U.S. Geological Survey Open-File Report 76-324, 77 p. *See Al-Garni 1996.* **(C4)**
- Hawley, J.W., compiler, 1978, Guidebook to the Rio Grande rift in New Mexico and Colorado: New Mexico Bureau of Mines and Mineral Resources, Circular 163, 241 p. **(A2, C2a, C4)**
- Clemons, R.E., 1979, Geology of Good Sight Mountains and Uvas Valley, southwest New Mexico: New Mexico Bureau of Mines and Mineral Resources, Circular 169, 31 p. *First map of Lake Goodsight* **(C2a, H1, I2)**
- Seager, W.R., and Morgan, P., 1979, Rio Grande rift in southern New Mexico, west Texas and northern Chihuahua, *in* Riecker, R.E., ed., Rio Grande rift, tectonics and magmatism: Washington D.C., American Geophysical Union, p. 87-106. **(C2a, C4)**
- Gates, J.S., White, D.E., Stanley, W.D., and Ackermann, H.D., 1980, Availability of fresh and slightly saline ground water in the basins of westernmost Texas: Texas Department of Water Resources, Report 256, 108 p. [First published as USGS Open-File Report 78-663, 115 p.] **(E2a, H2)**

C7.2.3. 1981 to 1995

- Gile, L.H., Hawley, J.W., and Grossman, R.B., 1981, Soils and geomorphology in the Basin Range area of southern New Mexico – Guidebook to the Desert Project: New Mexico Bureau of Mines and Mineral Resources, Memoir 39, 222 p. <https://geoinfo.nmt.edu/publications/monographs/memoirs/39/> **(C2a, C3)**
- Seager, W.R., 1981, Geology of the Organ Mountains and southern San Andres Mountains, New Mexico: New Mexico Bureau of Mines and Mineral Resources, Memoir 36, 97 p. **(C2a)**
- Wilson, C.A., White, R.R., Orr, B.R., and Roybal, R.G., 1981, Water resources of the Rincon and Mesilla Valleys and adjacent areas, New Mexico: New Mexico State Engineer Technical Report 43, 514 p. **(H1, H2)**
- INEGI, 1983c-cj, Ciudad Juárez H13-1, Cartas Hidrológica de Aguas: Instituto Nacional de Estadística, Geografía e Informática, SPP Programación y Presupuesto. Dirección General de Geografía del Territorio Nacional. Escala 1:250,000. **(F3)**
- Gates, J.S., White, D.E., and Leggat, E.R., 1984, Preliminary study of aquifers of the lower Mesilla Valley in Texas and New Mexico by model simulation: U.S. Geological Survey Water Resources Investigations Report 84-4317, 21 p. **(H1, H3)**
- Seager, W.R., Shafiqullah, M., Hawley, J.W., and Marvin, R.F., 1984, New K-Ar dates from basalts and the evolution of the southern Rio Grande: Geological Society of America Bulletin, v. 95, no. 1, p. 87-99. **(C2a, I3)**
- Wilson, C.A., and White, R.R., 1984, Geohydrology of the central Mesilla Valley, Doña Ana County, New Mexico: U.S. Geological Survey Water Resources Investigations Report 82-444, 144 p. **(H1, H2)**
- Orr, B.R., and White R.R., 1985, Selected hydrologic data from the northern part of the Hueco Bolson, New Mexico and Texas: U.S. Geological Survey Open-File Report 85-696, 88 p. **(H1)**
- Myers, R.G., and Orr, B.R., 1986, Geohydrology of the aquifer in the Santa Fe Group, northern West Mesa of the Mesilla Basin near Las Cruces, New Mexico: U.S. Geological Survey Water-Resources Investigations Report 84-4190, 37 p. **(C2a, H1)**
- Seager, W.R., Hawley, J.W., Kottowski, F.E., and Kelley, S.A., 1987, Geologic map of east half of Las Cruces and northeast El Paso 1° x 2° sheets, New Mexico: New Mexico Bureau of Mines and Mineral Resources, Geologic Map, GM-57, scale 1:125,000, 3 sheets. **(C2a)**
- Peterson, D.M., and Wilson, J.L., 1988, Variably saturated flow between streams and aquifers: New Mexico Water Resources Research Institute Report No. 233, 289 p. <https://nmwrri.nmsu.edu/publications/technical-reports/tr-reports/tr-233.html> **(D2, H3)**
- Bedinger, M.S., Sargent, K.A., and Langer, W.H., eds., 1989a, Studies of geology and hydrology in the Basin and Range province, southwestern United States, for isolation of high-level radioactive waste – Characterization of the Trans-Pecos region, Texas, U.S. Geological Survey Professional Paper 1370-B, 43 p. **(F2, H3)**
- Bedinger, M.S., Sargent, K.A., and Langer, W.H., eds., 1989b, Studies of geology and hydrology in the Basin and Range province, southwestern United States, for isolation of high-level radioactive waste – Characterization of the Rio Grande region, New Mexico and Texas: U.S. Geological Survey Professional Paper 1370-C. 42 p. **(F2, H3)**
- Bedinger, M.S., Sargent, K.A., and Langer, W.H., 1989c, Studies of geology and hydrology in the Basin and Range province, southwestern United States, for isolation of high-level radioactive waste – evaluation of regions: U.S. Geological Survey Professional Paper 1370-H, 61 p. **(D1)**

- Seager, W.R., 1989, Geology beneath and around the West Portrillo basalts, Doña Ana and Luna Counties, New Mexico: *New Mexico Geology*, v. 11, no. 3, p. 53-59. **(C2a)**
- Frenzel, P.F., and Kaehler, C.A., 1992, Geohydrology and simulation of ground-water flow in the Mesilla Basin, Doña Ana County, New Mexico and El Paso County, Texas, *with a section on Water quality and geochemistry by S.K. Anderholm*: U.S. Geological Survey Professional Paper 1407-C, 105 p. **(F2, H2, H3)**
- Hawley, J.W., and Lozinsky, R.P., 1992, Hydrogeologic framework of the Mesilla Basin in New Mexico and western Texas: New Mexico Bureau of Mines and Mineral Resources, Open-File Report 323, 55 p. **(H1)**
- Orr, B.R., and Risser, D.W., 1992, Geohydrology and potential effects of development of freshwater resources in the northern part of the Hueco Bolson, Doña Ana and Otero Counties, New Mexico, and El Paso County, Texas: U.S. Geological Survey Water-Resources Investigations Report 91-4082, 92 p. **(H1, H2)**
- Clemons, R.E., 1993, Petrographic analysis of Cenozoic-Mesozoic-Permian well cuttings from two exploration wells in south-central New Mexico: New Mexico Bureau of Mines and Mineral Resources, Circular 203, 28 p. **(C2b, H1)**
- Nickerson, E.L., and Myers, R.G., 1993, Geohydrology of the Mesilla ground-water basin, Doña Ana County, New Mexico, and El Paso County, Texas: U.S. Geological Survey Water-Resources Investigations Report 92-4156, 89 p. **(H1, H2)**
- Hanson, R.T., McLean, J.S., and Miller, R.S., 1994, Hydrogeologic framework and preliminary simulation of ground-water flow in the Mimbres Basin, southwestern New Mexico: U.S. Geological Survey Water-Resources Investigations Report 94-4011, 118 p. **(F2, H1, H3)**
- Seager, W.R., 1995, Geology of southwest quarter of Las Cruces and northwest El Paso 1° x 2° sheets: New Mexico Bureau of Mines and Mineral Resources, Geologic Map 60, scale 1:125,000. **(C2b, C4)**

C7.2.4. 1996 to 2010

- Woodward, D.G., and Myers, R.G., 1997, Seismic investigation of the buried horst between the Jornada del Muerto and Mesilla ground-water basins near Las Cruces, Doña Ana County, New Mexico: U.S. Geological Survey Water-Resources Investigations Report 97-4147, 45 p. **(C4, H1)**
- Hibbs, B.J., Creel, B.J., Boghici, R., Hayes, M., Ashworth, J., Hanson, A., Samani, Z., Kennedy, J.F., Hann, P., and Stevens, K., 1997, Transboundary Aquifers of the El Paso/Ciudad Juárez/Las Cruces Region: U.S. Environmental Protection Agency, Region 6; Technical Contract Report- Interagency Contracts X-996343-01-0 and X-996350-01-0, prepared by the Texas Water Development Board and the New Mexico Water Resources Research Institute, variously paged. *See TWDB, 1997, Appendix C—G.I.S. coverages, metadata descriptions, [and] groundwater data sets on CD-ROM, with Water Quality map insert on back-cover.* <https://nmwrri.nmsu.edu/publications/publications.html> **(F1, H1, H2)**
- Texas Water Development Board (TWDB), 1997, Appendix C – G.I.S. coverages, metadata descriptions,[and] groundwater data sets on CD-ROM, with Water Quality map* insert on back-cover; *in* Hibbs and 9 others, Transboundary Aquifers of the El Paso/Ciudad Juárez/Las Cruces Region: U.S. Environmental Protection Agency, Region 6; Technical Contract Report- Interagency Contracts X-996343-01-0 and X-996350-01-0, prepared by the Texas Water Development Board and the New Mexico Water Resources Research Institute, variously paged. **Map includes Stiff diagrams color-coded for four total dissolved solids classes [0-1,000, 1,000-3,000, 3,000-5,000, and >5,000 mg/L] for more than 200 wells in Doña Ana and Otero Counties, NM; El Paso and Hudspeth Counties, TX; and contiguous parts of Chihuahua.* <https://nmwrri.nmsu.edu/publications/publications.html> **(F1, H2)**
- Hibbs, B., Boghici, R., Ashworth, J., Hayes, M., Peckham, D., Guillen, R., Fuentes, O., Laloith, N., Morales, M., Maldonado, A., Creel, B., Kennedy, J., Hanson, A., Samani, Z., Nunez, F., Lemus, R., Moreno, G., Rascon, E., Kuo, R., Waggoner, S., Ito, C., Robinson, J., Valdez, J., Little, D., Rascon, A., Reyes, A., Williams, K., Vaughan, M., Cabra, O., Kelly, T., and King, C., 1998, Transboundary aquifers and binational ground-water data base, City of El Paso/Ciudad Juárez area; Base de datos binacional del acuífero transfronterizo, de Ciudad Juárez, Chih./El Paso, Tex.: first binational aquifer report and data base sanctioned by the governments of the United States and Mexico, 47 p. + appendices and CD-ROM. Participating agencies; International Boundary and Water Commission, U.S. Environmental Protection Agency, Texas Water Development Board, New Mexico Water Resources Research Institute, Comisión Internacional de Límites y Aguas, Comisión Nacional del Agua, Junta Municipal de Agua y Saneamiento de Ciudad Juárez. **(F1, H1, H2)**

- Nickerson, E.L., 1998, U.S. Geological Survey seepage investigations of the Lower Rio Grande in the Mesilla Valley, *in* Ortega Klett, C.T., ed., Proceedings of the 43rd Annual New Mexico Water Conference: Water Challenges on the Lower Rio Grande. New Mexico Water Resources Research Institute Report No. 310, p. 59-68. **(H3)**
- Hawley, J.W., and Kernodle, J.M., 2000, Overview of the hydrogeology and geohydrology of the northern Rio Grande basin – Colorado, New Mexico, and Texas, *in* Ortega Klett, C.T., ed., Proceedings of the 44th Annual New Mexico Water Conference: New Mexico Water Resources Research Institute Report No. 312, p. 79-102. **(D1)**
- Hawley, J.W., Hibbs, B.J., Kennedy, J.F., Creel, B.J., Remmenga, M.D., Johnson, M., Lee, M.M., and Dinterman, P., 2000, Trans-International Boundary aquifers in southwestern New Mexico: New Mexico Water Resources Research Institute, prepared for U.S. Environmental Protection Agency-Region 6 and International Boundary and Water Commission; Technical Completion Report-Interagency Contract X-996350-01-3, 126 p. <https://nmwrri.nmsu.edu/publications/pub-external-pages/trans-international-boundary-aquifers-in-southwest-new-mexico.html> **(C2b, D1, F1, I2)**
- Jiménez, A.J., and Keller, G.R., 2000, Rift basin structure in the border region of northwestern Chihuahua: New Mexico Geological Society Guidebook 51, p. 79-83. **(C2b, C4, F1, H1)**
- Kennedy, J.F., Hawley, J.W., and Johnson, M., 2000, The hydrogeologic framework of basin-fill aquifers and associated ground-water-flow systems in southwestern New Mexico – An overview: New Mexico Geological Society Guidebook 51, p. 235-244. **(F1, H1, I2)**
- Phillips, F.M., Hogan, J.F., Mills, S.K., and Hendrickx, J.M.H., 2003, Environmental tracers applied to quantifying causes of salinity in arid-region rivers: Preliminary results from the Rio Grande, southwestern USA, *in* Alsharhan, A.S., and Wood, W.W., eds., Water resources perspectives: Evaluation, management and policy: Amsterdam, Elsevier, p. 327-334. **(D2, H2)**
- Hawley, J.W., and Kennedy, J.F., 2004, Creation of a digital hydrogeologic framework model of the Mesilla Basin and southern Jornada del Muerto Basin: New Mexico Water Resources Research Institute Report No. 332, 105 p., with plates and appendix on CD ROM. <https://nmwrri.nmsu.edu/publications/technical-reports/tr-reports/tr-332.html> **(F1, H1)**
- Witcher, J.C., King, J.P., Hawley, J.W., Kennedy, J.F., Williams, J., Cleary, M., and Bothern, L., 2004, Sources of Salinity in the Rio Grande and Mesilla Basin Groundwater: New Mexico Water Resources Research Institute Report No. 330, 168 p., with appendices and plates on CD ROM. <https://nmwrri.nmsu.edu/publications/technical-reports/tr-reports/tr-330.html> **(C2b, E2a, H1, H2)**
- Connell, S.D., Hawley, J.W., and Love, D.W., 2005, Late Cenozoic drainage development in the southeastern Basin and Range of New Mexico, southeasternmost Arizona and western Texas, *in* Lucas, S.G., Morgan, G., and Zeigler, K.E., eds., 2005, New Mexico's Ice Ages: New Mexico Museum of Natural History & Science Bulletin No. 28, p. 125-150. **(C2b, I1, I3)**
- Hawley, J.W., Kennedy, J.F., Ortiz, M.A., and Carrasco, S., 2005, Creation of a digital hydrogeologic framework model of the Rincon Valley and adjacent areas of Dona Ana, Sierra and Luna Counties NM: New Mexico Water Resources Research Institute, Addendum to Report No. 332 on CD ROM. <https://nmwrri.nmsu.edu/publications/technical-reports/tr-reports/tr-332.html> **(H1)**
- Hibbs, B.J., and Darling, B.K., 2005, Revisiting a classification scheme for U.S.-Mexico alluvial basin-fill aquifers: Ground Water, v. 43, no. 5, p. 750-763. **(D1, D2, F1)**
- Hutchison, W.R., 2006, Groundwater management in El Paso, Texas: University of Texas at El Paso, Center for Environmental Resource Management, doctoral dissertation, 329 p. **(H1, H2, H3)**
- Mack, G.H., Seager, W.R., Leeder, M.R., Perea-Arlucea, M., and Salyards, S.L., 2006, Pliocene and Quaternary history of the Rio Grande, the axial river of the southern Rio Grande rift, New Mexico, USA: Earth-Science Reviews, v. 77, p. 141-162. **(C2b, I3)**
- Nickerson, E.L., 2006, Description of piezometers and ground-water-quality characteristics at three new sites in the Lower Mesilla Valley, Texas, and New Mexico: U.S. Geological Survey Scientific Investigations Report 2005-5248, 27 p. **(H1, H2)**
- Hawley, J.W., and Kernodle, J.M., 2008, Early contributions to arid-zone hydrogeology in the eastern Basin and Range region: Ground Water, v. 46, no. 3, p. 510-516. **(A2, B3, D1)**
- Hawley, J.W., Kennedy, J.F., Granados Olivas, A., and Ortiz, M.A., 2009, Hydrogeologic framework of the binational western Hueco Bolson-Paso del Norte area, Texas, New Mexico, and Chihuahua: Overview and progress report on digital model development: New Mexico Water Resources Research Institute Report No. 349, 45 p., 2 pls. <https://nmwrri.nmsu.edu/publications/technical-reports/tr-reports/tr-349.html> **(F1, H1)**

C7.3. Topic/Subtopic Categories, with Alphanumeric Cross-Reference Codes (Appendix B)

A. Bibliographies, Dictionaries, Glossaries, Biographies, Reviews, and News Items

- A1. Bibliographies, Dictionaries, and Glossaries
- A2. Biographies and Reviews
- A3. News Items

B. Time: Geologic, Prehistoric, and Historic

- B1. Geologic and Prehistoric Time
- B2. Prehistoric Perspective: US Southwest and Northern Mexico
- B3. Historic Perspective: US Southwest and Northern Mexico

C. Environmental, Physiographic, and Geologic Setting

- C1. Climatic, Hydrographic, Ecologic, and Paleoenvironmental Setting
- C2. Geologic and Geomorphic Setting
 - C2a. Geologic and Geomorphic Setting: Pre-1990
 - C2b. Geologic and Geomorphic Setting: Post-1989
- C3. Soil-Geomorphic Relationships and Soil Surveys
- C4. Geophysical/Geochemical Data and Interpretations

D. Basic Hydrogeologic Concepts

- D1. Conceptual Models, Definitions, and Regional Overviews
- D2. Groundwater-Flow Systems, Including Recharge

E. GIS/Remote Sensing and GW-Resource Management/Planning

- E1. GIS/Remote Sensing
- E2. Resource Management/Planning
 - E2a. Desalination
 - E2b. Recharge and Recovery
 - E2c. Groundwater-Quality Projection and Waste Management
- E3. Legal and Environmental Issues and Constraints

F. Transboundary Regional Hydrogeology and Geohydrology

- F1. Binational
- F2. USA
- F3. México

G. Early Documents on Mesilla Basin Regional Aquifer Systems (1858-1970)

- G1. 1858 to 1935
- G2. 1935 to 1970

H. Contemporary Documents on Mesilla Basin Regional Aquifer Systems

- H1. Hydrogeology
- H2. Hydrochemistry
- H3. Flow Models

I. Paleohydrology: Ancestral Fluvial and Pluvial Lake Systems

- I1. Regional Overviews
- I2. Transboundary Region Paleohydrologic Systems
- I3. Evolution of the Rio Grande Fluvial System

C8a. SPECIAL ADDENDUM 1—HAWLEY, J.W., 1971, GEOLOGY AND ITS RELATION TO THE HYDROLOGIC SYSTEM (Reformatted, with Original Pagination inserts) * **

**Contribution approved for publication by the Director, Soil Conservation Service, U.S. Department of Agriculture.*

***Pages 9 to 24 in King, W.E. and others, 1971, Geology and ground-water resources of central and western Doña Ana County, New Mexico: N.M. Bureau of Mines and Mineral Resources, Hydrologic Report 1.*

[p. 9] GEOMORPHIC FEATURES

The area under discussion (figs. 1, 3) is in the Mexican Highland section of the Basin and Range physiographic province (Fenneman, 1931; Thornbury, 1965). It includes a number of mountain uplifts, parts of four major intermontane basins, and the valley of the Rio Grande. The area is bounded on the west by the Mimbres River basin, the Good sight Mountains, and the Hillsboro-Animas uplift. The San Andres-Organ-Franklin Mountain chain forms the eastern boundary. The intermontane basins extend southward into northern Chihuahua and northward into central New Mexico.

The geomorphology of the region has been discussed in detail by Ruhe (1962, 1964, 1967), Kottowski (1958), Kelley and Silver (1952), and Hawley (1965; Hawley and Gile, 1966; Hawley and Kottowski, 1969). The succeeding discussion will emphasize features considered to be of hydrogeologic significance.

MOUNTAINS

With some major exceptions, the mountains in the Rio Grande region of southern New Mexico consist of fault-block uplifts with a general north-south trend (Kottowski, 1958). Other mountain types include broad domal uplifts and remnants of igneous intrusive bodies. Only the low-lying West Potrillo Mountains (maximum elevation, 5,473 ft), in south-west Doña Ana County, reflect the primary form of the basaltic volcanic cones and flows that underlie that particular upland area.

The highest peak in the region, Organ Needle (elevation, 9,012 ft) is part of the jagged crest of a mass of Tertiary [igneous] intrusive rocks (monzonites) that comprises the central Organ Mountains. The San Andres range and the Franklin and southern Organ Mountains generally have the form of strongly tilted fault blocks, which are bounded on the west by dip slopes on Paleozoic to Tertiary beds (dominantly marine carbonate rocks), and on the east by escarpments that cut across Paleozoic strata into Precambrian rocks. Alluvial- fan and pediment gravels on the upper piedmont slopes of the western Hueco and Tularosa Basins are cut by faults that are closely aligned with the eastward-facing escarpments. The Caballo Mountains, which extend into the north-central part of the area of investigation, are similar in form to the San Andres-Franklin range, with the exception that they consist of several eastward-tilted fault blocks.

The Robledo Mountains also consist of a tilted fault-block uplift, but in this case, the mountains have the form of a wedge-shaped horst that is bounded on the east and west by faults and is tilted gently toward the south. The peaks and high ridges of the Robledo, Caballo, San Andres, and Franklin Mountains are in most places underlain by thick-bedded carbonate rocks of Paleozoic age.

The two other major upland areas, the Doña Ana Mountains and the Sierra de Las Uvas, are domal uplifts composed mainly of Tertiary igneous rocks. Monzonite intrusive rocks form the high peaks of the Doña Ana Mountains, and thick basaltic andesite flows and rhyolitic welded tuffs hold up high mesas, cuestas, and buttes in the Sierra de Las Uvas. The west flanks of the Uvas uplift make up the east limb of a large northward-plunging syncline. The Tertiary volcanics cropping out in the nose and west limb of this syncline form the arcuate Good sight Mountains.

A number of small, isolated upland areas occur throughout the region. Many are aligned along definite trends and appear to be associated with buried extensions of major positive structures. Small fault-block uplifts include the Rincon Hills, Tonuco-Selden Hills uplift, Tortugas Mountain, and Bishop Cap. Small peaks formed by erosional remnants of Tertiary igneous intrusive bodies include Goat Mountain, Picacho Peak, Vado Hill, Mount Riley, and Cerro de Muleros [Cristo Rey]. The Aden, Sleeping Lady, and Rough and Ready Hills consist of a belt of small peaks, ridges, buttes, and elongated mesas underlain by Tertiary volcanic rocks. The hills appear to be remnants of a former cover of andesites, basalts, rhyolite tuffs, and associated sedimentary rocks that had been disrupted by a combination of erosion, faulting, and warping. Point of Rocks, between the southern Caballo and San Andres ranges, is another group of hills of this type.

**C8b. SPECIAL ADDENDUM 2—PROJECT PROPOSAL FOR A COMPREHENSIVE STUDY
OF THE WATER RESOURCES OF THE SOUTHERN RIO GRANDE VALLEY AREA,
NEW MEXICO (C.A. Wilson, May 1972; Reformatted, with original pagination inserts)**

PROJECT PROPOSAL FOR A COMPREHENSIVE STUDY OF THE WATER RESOURCES OF THE
SOUTHERN RIO GRANDE VALLEY AREA, NEW MEXICO

C.A. Wilson

UNITED STATES DEPARTMENT OF THE INTERIOR, GEOLOGICAL SURVEY
Albuquerque, New Mexico, May 1972

CONTENTS

INTRODUCTION -----	4
NEED FOR THE STUDY -----	8
PREVIOUS STUDIES AND DATA AVAILABILITY -----	13
OBJECTIVES AND PLAN OF STUDY -----	18
Water resources in vicinity of Las Cruces -----	22
Water resources in the Rincon and Mesilla Valleys -----	24
Major items of work -----	30
Schedule of work, time, and costs -----	34
Reports -----	40
SELECTED REFERENCES -----	42


ILLUSTRATIONS

Figure 1. Map of lower Rio Grande Valley area, New Mexico -----	5
---	---

INTRODUCTION [p. 4-7]

The water supply of the lower Rio Grande Valley and adjacent uplands is derived from surface water from the Rio Grande and ground water from sediments of the intermontane basin. Neglecting the short-term effects the surface and ground-water units function as one system due to hydraulic connection between the river and the aquifers. Planning the orderly development of the water resources of the valley requires current factual data followed by complete and reliable analyses.

The purpose of this proposal is to aid in the planning, organization, and preparation of a comprehensive study plan of the water resources of the southern Rio Grande Valley in New Mexico, to evaluate data presently available, and to determine additional data that will need to be collected.



[p. 5] The following agencies might be cooperators with the U.S. Geological Survey for the proposed study:

State Engineer of New Mexico
City of Las Cruces
Elephant Butte Irrigation District
New Mexico Water Resources Research Institute

The New Mexico Water Resources Research Institute has furnished office space to the Geological Survey since the project began in October 1971 and has extended this privilege for the duration of the project.

The lower Rio Grande Valley is in the basin and range province of southwestern New Mexico, mostly in Dona Ana and Sierra Counties (fig. 1 [p.6]). The western boundary of the basin is formed by the East Potrillo, West Potrillo, Goodsight Mountains, and the Black Range; the eastern boundary by the Franklin, Organ, and San Andres Mountains.

The Rio Grande Irrigation Project of the Bureau of Reclamation encompasses the river and adjacent flood plain downstream from Caballo Reservoir to below EL Paso, Tex. The upper part is

referred to as the Rincon Valley; it extends from Caballo Reservoir, in Sierra County, southward to the narrowing of the flood plain, near Tonuco in Dona Ana County. The Mesilla Valley extends from near Radium Springs, Dona Ana County, to the "narrows" at El Paso. About 180,000 acres are served with irrigation water by the Project. About 100,000 irrigated acres are in Rincon and Mesilla Valleys.

[p. 7] The largest use of water within the valley is for irrigation (mostly surface water with supplemental supplies from ground water). During periods of drought, ground water is the major source of irrigation water. Ground water is the municipal supply, for Las Cruces Anthony, Hatch, and most of the supply (over 85 percent) for El Paso. Most rural areas use ground water for domestic and stock needs; and industry depends heavily on ground-water supplies.

On May 4, 1971 the Governors of New Mexico and Texas gave support to a proposed areal resource study, the "Rio Grande Regional Environmental Study." Elephant Butte Reservoir, New Mexico--Fort Quitman, Texas. The overall purpose of this large study is to enhance 'the social, economic, and environmental conditions of the valley, through inventory, study, and planned use of the total resources available to the area, especially the surface and ground-water supplies. The study should evaluate resource development and management plans using multipurpose approaches to the resource problems. The major resource problems to be dealt with by the Environmental Study are water resources and supply.

NEED FOR THE STUDY [p. 8-12]

The needs for a study of the water resources in the southern Rio Grande Valley, New Mexico may be categorized into three general Groups:

- (1) the need for information on the general hydrology over the entire area, with emphasis on the occurrence, availability and quality of the ground water,
- (2) greater detailed information is needed by the city of Las Cruces for planning and developing municipal supplies for the future,
- (3) evaluation of the availability, quantity, quality, and uses of ground water in the flood-plain alluvium, and adjacent units along the Rio Grande, and recommendations or considerations in the design and operation of combined ground- and surface-water system for year-round irrigation

[p. 9] Recent studies have defined the geologic framework for the occurrence of ground water in the Rio Grande Valley (see References). These studies need to be expanded and knowledge increased on the general hydrologic situation, especially the availability, quantity, and chemical quality of the ground water, and the aquifer characteristics of the intermontane valley fill. In many areas of the basin, basic data on ground water are scarce. Some investigators feel that large quantities of potable water are present in La Mesa bolson; others believe the quantity is limited. Little basic data are available on ground water in the southern part of the Jornada del Muerto. The connection, if any, between the aquifers which furnish water to Las Cruces and to El Paso's Canutillo well field is known. The connection and extent of subsurface boundaries need to be defined throughout the basin, little is known about ground-water conditions below depths of a few hundred feet. It is felt that additional hydrologic studies would give information and may provide answers to these problems.

A port of entry from Mexico into New Mexico is being proposed for Anapra, located about 10 miles west of downtown El Paso. This gateway to Mexico would cause industrial and agricultural requirements for water to increase greatly in that area. However, ground-water conditions are not known in the vicinity of Anapra. Water resources should be determined during the earliest planning phase of planning the port of entry.

[p. 10] The city of Las Cruces has developed its water supply from about 18 wells finished in sand and gravel units of the Santa Fe Group (one of these wells taps the Rio Grande flood-plain alluvium). The wells are located within the city limits, some exceeding 750 ft in depth. Some increase in the dissolved-solids content of the water has been noticed in recent years. Also, recent maps showing the altitude of the water table in the vicinity of Las Cruces indicate that ground water of higher dissolved-solids content may be moving downgradient from the river and flood-plain alluvium into the cone of depression created by pumping wells in the city well field.

King, and others (1969, p. 18-20) discuss the subsurface occurrence of nearly impermeable igneous intrusive barriers to ground-water movement in the vicinity of the well field. These intrusives tend to limit expansion into adjoining areas without detailed exploration to insure that new wells will encounter sufficient saturated thickness and permeable sediments.

City officials need detailed information on the extent and volume of fresh-water supplies in the vicinity of Las Cruces, delineation of areas where new wells should be drilled, and recommendations for development so as to recover the fresh water most effectively. Also, the city needs to know what changes in chemical quality may occur in the future due to pumpage. Development of the fresh-water zones must take into account and suggest ways to prevent or limit the possible encroachment of underlying and/or overlying more saline water.

[p. 11] Irrigation uses and reuses the largest amount of water in the Rincon and Mesilla Valleys. Often the supply of surface water in the river is deficient, and supplemental irrigation using ground water from the adjacent shallow (± 80 feet) flood-plain alluvium is needed. The ground water pumped is replenished later during subsequent irrigation by surface water in years of above-normal supply. Early hydrologic studies of this irrigation system indicated that any substantial ground-water pumpage would result in much lower drain return flow and hence less availability of surface water downstream in the next 3 or 4 years. However, ground-water pumpage during the droughts of the early and mid-1950s and again in the mid-1960s did not appear to cause the predicted later severe decreases in surface-water supplies downstream. Overall, the effect of ground-water pumpage on the river needs further study. Quantities in the field need to be measured as much of the early work was based on estimated values and/or calculation.

The chemical quality of the water in the alluvium is variable; in general, the water is more highly mineralized than water in the river, or in the upper aquifers of the adjacent Santa Fe Group. This mineralization is probably caused by use and reuse of water for irrigation. Definition is needed on the quality of this shallow alluvium water, its circulation patterns, the depth to which fresher and more saline water may occur in the underlying aquifers, and changes in water-quality which occur. Sources of possible water pollution need to be delineated and monitored.

[p. 12] Basic needs other than to the surface-ground water irrigation system in the lower Rio Grande Valley would include, (1) the capability for year-round irrigation from a firm supply of water; (2) improvement in the quality of irrigation water applied in the fields, and improvement in the quantity and quality of return flow which is used to meet downstream demands; and (3) the reduction of water losses in the current delivery system. In order to meet these needs, the Elephant Butte Irrigation District is planning revisions to the physical layout and construction of the irrigation network that will involve a combined surface-ground water system. These revisions will require hydrologic information to aid in design and modification of facilities, and to allow predictions of water quality and quantity changes due to different demands or stresses which may be applied in the course of managing the irrigation system. The ground-water reservoir must be defined in order to best meet these needs and techniques developed to allow management of the whole hydrologic system.

In addition to the southern Rio Grande Valley area, New Mexico, the Mimbres River Basin and Tularosa Valley areas may be of importance to the Regional Environmental Project. At present, water-resources studies are being done in these areas, and information from the studies are available.

PREVIOUS STUDIES AND DATA AVAILABILITY [p. 13-17]

Many geologic and hydrologic studies have been made in the lower Rio Grande Valley since the first work in 1905. Many references to these published studies and some open-file reports of various governmental agencies are included in the Reference section of this proposal.

The first ground-water study in the valley was by Slichter (1905). By use of observation wells he demonstrated the connection between the river and the alluvium aquifer, and calculated the underflow of the river at the "narrows" near El Paso. He conducted some aquifer tests on wells and described well construction and pump design.

Keyes (1905) recorded the geology and ground-water conditions in the Jornada del Muerto. The report is elaborate in its descriptions of the desolation of this area. Keyes subdivided the area into general geologic units and outlined structural relations. Data are given on the occurrence of ground water and some wells are described.

Lee (1907) reported on the water resources of the entire Rio Grande Valley of New Mexico. He noted the large discharge of ground water by evaporation in the valley and speculated that the flow of ground water in La Mesa was toward the Rio Grande.

The geology of Dona Ana County was studied by Dunham (1935). He recognized the basin-fill deposits as belonging to the Santa Fe Group and outlined the formation of the Mesilla Valley.

[p. 14] The most extensive ground-water investigation in the entire basin area was by Conover (1954). The main purpose was to study possibilities of pumping ground water for supplemental irrigation when surface-water supplies were deficient, and the effects of this pumpage on later flow in drains and the river. Conover concluded, through use of gross water-budget calculations, that only a fixed amount of water was available in the Rincon and Mesilla Valleys, and that heavy ground-water pumpage over a period of several years would cause less return flows to be available for reuse downstream by other irrigators in the succeeding 3 to 4 years because of replacing the water pumped from storage. His report contains records for 108 wells in the flood-plain area, 147 wells on the uplands, 44 chemical analyses of well water, and 45 driller's logs. He estimated that the river alluvium had an average transmissivity of about 75,000 gpd/ft (gallons per day per foot) and that the upland parts of the Santa Fe Group had an average transmissivity of about 30,000 gpd/ft. These estimates were based on seven aquifer tests and seven calculations of transmissivity based on drain discharges and water-table gradients. His report also contains a map showing the altitude of the water table over most of the basin area.

Gunaji (written communication 1961) differed with some previous investigators on the effect of ground-water pumpage on return flows into the river. He showed that aquifer recover from the heavy ground-water pumpage of 1954-1956 was more rapid than others would have predicted. He felt that ground water should be used as supplemental supply during years of short surface supplies. Gunaji's report contains no additional field data that is not in previous reports.

[p. 15] Three aquifers (shallow, medium, and deep) were described by Leggat, Lowry, and Hood (1962) in the Canutillo well field south of Anthony. Detailed data on the lower Mesilla Valley area of New Mexico and Texas shows leakage between the three aquifers and changes in flow patterns due to changes in pumpage from the different aquifers. Their report showed the quick recovery in 1958 of water levels in the alluvial aquifer from previous heavy pumpage during years of drought. Occurrences of fresh water at depths over 1,200 feet were noted. Some estimates on water-availability are calculated. The report contains records of 219 wells and 76 driller's logs, and about 350 chemical analyses of water; most wells and test holes are located in Texas.

Hawley (1965), and Hawley and Gile (1966) discuss the geomorphic surfaces and the recent geologic formation of the lower Rio Grande Valley area. These reports are quite detailed and show the chronological development of the basin and deposition of the Santa Fe Group from Paleocene to the present.

The most recent hydrogeology investigation is by King, Hawley, Taylor and Wilson (1969). This report pertains to ground-water conditions in the Santa Fe Group and includes a map showing the altitude of the water table throughout much of the basin. The report describes and relates the occurrence of ground water to geology. The report contains records for 219 wells and test holes, 39 chemical analyses, and driller's logs or lithologic descriptions of samples from 52 wells and test holes. Some of these well records are updated measurements of wells listed by Conover (1964).

[p. 16] Some unpublished sources of hydrologic information on the lower Rio Grande Valley include reports by Basler and Alary (1968), Doty (1969), Meyer and Gordon (1971), and Richardson (1971). The purpose of the report by Basler and Alary was to determine the quality of the shallow ground water in the alluvium of the flood plain in Rincon and Mesilla as an aid in planning a comprehensive study of ground water in the valley of the Rio Grande. Forty wells were sampled by Basler and Alary. Doty (1969) reported on development of the water supply for the NASA site in eastern Dona Ana County. Meyer and Gordon (1971) is the most current progress report (covers the period 1963-70) on the development of ground water in the El Paso district, Texas. The report updates information on development, well locations, and pumpage; it also discusses the rapid recovery of ground-water levels in the alluvium after the heavy pumpage of 1964. The report contains the water budget for 1968 of the lower valley area (the Rio Grande Valley below El Paso). Inflow from surface water, ground water, and precipitation amounted to 348,700 acre-feet; outflow as surface-water flow, ground-water movement, consumptive use, evaporation, and changes in storage amounted to 314,100 acre-feet. The imbalance, about 10 percent or 34,600 acre-feet, may represent such things as use or flow into Mexico, errors in measurements or estimates, and pumpage out of the river.

[p. 17] Richardson (1971) constructed a mathematical model of the Mesilla Valley based on a model program written by Dr. William Brutsaert of New Mexico Institute of Mining and Technology. Basic data used in the model were taken from published and unpublished sources. Model calibration was by changing the value of the storage coefficient (ranged from 15 to 25 percent best fit using 20 percent). Model verification was by comparing the average of monthly water-level measurements for 1962 and 1964 in 21 selected observation wells, with the average of computed water-level values in the same grid nodes as the observation wells.

Some data are available in the files of various government agencies, however, much may have been published. The Geological Survey files (New Mexico District) for Dona Ana County contain about 350 well Schedules, 225 chemical analyses (some wells may have several analyses), 8 borehole geophysical logs, 7 aquifer-test results, and 65 driller's logs. The Geological Survey files (Texas District) contains about 250 well schedules for wells located below Anthony in the lower Mesilla Valley of Texas and New Mexico. About 170 well schedules, 150 chemical analyses, no borehole geophysical logs, 38 aquifer tests, and 10 driller's logs are available for Sierra County.

Records of stream flow and flow in canals and drains are available in published and unpublished records of the Bureau of Reclamation, International Boundary and Water Commission, Elephant Butte Irrigation District, and the U.S. Geological Survey.

OBJECTIVES AND PLAN OF STUDY [p.18-21]

The boundaries of the proposed water-resources study of the southern Rio Grande Valley are the latitude of Caballo Reservoir on the north, the San Andres, Organ, and Franklin Mountain ranges on the east, the "narrows" near El Paso, Tex., and the Mexican border on the south, and the Mimbres-Rio

Grande watershed boundary (approximates the East and West Potrillo, and Good sight Mountain ranges) on the west (fig. 1).

[p. 19] The objectives of the general hydrology study of the entire area would include, but not be limited to, the study and determination of the following:

- (1) thickness of the river flood-plain alluvium; thickness of the Santa Fe Group; delineation of major aquifers; and/or water-quality zones,
- (2) water quality and variations in quality with areal extent and depth, and if possible, the base of the fresh and slightly saline water zones (water containing 1,000 mg/l, milligrams per liter, or less and 1,000 to 3,000 mg/l respectively of dissolved solids),
- (3) the altitude of the water table and/or potentiometric surface, and the general flow patterns of the ground water in the different aquifers,
- (4) the thickness of the fresh and slightly saline water zones, and the total saturated thickness of the alluvium and the Santa Fe Group,
- (5) the amount of ground water in storage in the various aquifers or zones,
- (6) areas, sources, and estimates of the amount of recharge to, and discharge from, the various aquifer,
- (7) depth to water throughout the area,
- (8) aquifer properties, such as transmissivity, storage, capacity, lithology, and structure,
- (9) availability of fresh and slightly saline water, and estimates of the potential yield of wells.

[p. 20] A detailed and well-planned program of data acquisition is necessary to meet the objectives of the general hydrology study. Several deep test holes are needed (about 8 holes to depths of approximately 1,500 feet) at selected locations throughout the study area. Several types of borehole geophysical logs would be run in the holes. Four types of data could be collected at selected depth intervals in each hole: (1) water-quality samples, (2) aquifer permeability, (3) hydrostatic pressure, and (4) aquifer lithology. If possible, the test holes should be converted and equipped as permanent observation wells so that changes in water quality and hydrostatic pressure with time can be recorded.

Magnetic, gravity, and seismic geophysical surveys, borehole geophysical logs, and selected driller's logs would be used to define the thickness of the Santa Fe Group and aquifer zones within the group. Information on water-quality zones would be obtained from water sampling at selected depths in test holes and in some wells, interpretation of borehole geophysical logs, and possibly by use of earth-resistivity measurements. The top of the water table and potentiometric surface would be based on water-level measurements in wells and test holes. Flow pattern can be outlined from vertical flow-meter tests and by head measurements at selected depths in test hole. Surface geological mapping may show areas of natural recharge and aquifer lithology. Data on aquifer properties, ground-water availability, and potential yield would be obtained by making aquifer tests (pump tests involving drawdown-yield measurements) on wells and test holes. The results of the general hydrologic study would be prepared as a comprehensive report for publication by the State or by the Geological Survey.

[p. 21] Specific areas in the lower Rio Grande Valley may require more detailed hydrologic work. For example, a study of ground-water resources in the vicinity of the proposed port of entry at Anapra needs to be done in as much detail as possible. This study should include the drilling of test holes to investigate water quality and aquifer yields.

WATER RESOURCES IN THE VICINITY OF LAS CRUCES [p. 22-23]

In the vicinity of Las Cruces the study would be expanded and done in greater detail. Special attention would be given to delineation of geologic trends and the location of subsurface barriers to flow within the aquifers so that areas favorable for future exploration for fresh water can be outlined. The

relationship of the aquifers from which Las Cruces, Anthony, and El Paso (Canutillo well field) produce water would be investigated. The decline of the water table around Las Cruces may cause water of inferior quality that moves downgradient in the river alluvium to enter the aquifer tapped by city wells. The extent of this decline, potential chemical-quality problems in the well field, and recommendations to relieve the problems would be studied.

The city of Las Cruces has information on most of the city water wells. Additional information is available on wells at New Mexico State University and in the files of various governmental and private institutions. It is planned to drill two or three of the eight proposed deep test bores in the vicinity of Las Cruces and several shallower test holes in the area between the river alluvium and the city well field. Gravity and seismic surveys would be done in greater detail around the city so that subsurface features can be delineated. Borehole geophysical-logs could be made in used or abandoned wells and test holes.

[p. 23] An open-file report on the water resources in the vicinity of Las Cruces would be prepared upon completion of that part of the study. The detailed report will contain maps and/or discussion of objectives 1 through 9 of the general hydrology study of the entire area.

WATER RESOURCES IN THE RINCON AND MESILLA VALLEYS [p. 24-29]

The third area of need water-resources study involves the surface-ground water system of the Rio Grande and associated irrigation network, and ground water in the surrounding flood-plain alluvium. Broad general objectives required to fulfill the needs discussed previously would involve collecting and analyzing sufficient basic data to define the hydrologic system in the valley area and then using this information to construct models which can predict reaction to desired management practices.

Specific objectives sought in relation to defining the hydrologic system include:

- (1) Evaluation of the usefulness of the mathematical model constructed by Richardson (1971). This can be done with data now in the report, and should show if the model could form the basis for future use in design, operation and management of the system. The model may also indicate areas of concentration in collecting basic data.
- (2) Determine quantities and discuss components of the water budget of the river and ground-water system in the valleys. Water-budget amounts for the Mesilla Valley were calculated for 1962 and 1964 for use in Richardson's model. Refinement of budget factors is needed, especially where values such as streamflow or ground-water pumpage can be measured.
- (3) Based on field measurements, delineate in detail and use maps when possible, the following items for both the flood-plain alluvium and adjacent parts of the Santa Fe Group [p. 24]:
 - a. Altitude of the base of the alluvial aquifer and effective hydrologic boundaries of adjoining aquifers.
 - b. Altitude of the water table and/or potentiometric surface.
 - c. Saturated thickness of the alluvium and the saturated thickness in other aquifers that are affected by large-scale pumpage.
 - d. Degree and areal extent of hydraulic connection between aquifers.
 - e. Water-quality zones within the aquifers.
 - f. Aquifer properties such as transmissivity and storage capacity.
 - g. Amount of ground-water pumpage.
- (4) Discuss and illustrate the relationship, hydraulic connection, and flow pattern of the ground water in the alluvium and in the Santa Fe Group.
- (5) Determine areas and quantities of better quality water in the alluvium and in nearby parts of the Santa Fe Group. Factors affecting and controlling the quality should be found and related to changes which may have occurred in recent years in water quality.

[p.26] Once the hydrologic system is defined (items 1-5), the remaining, but most important objective is to determine future changes in quantities and chemical quality of the irrigation water due to management manipulation of the system. It is suggested that a digital or combination analog-digital model be constructed and used to meet this objective. Many recommendations for operation of the system can be made in the course of analyzing the basic data collected (such as areas and zones of the better quality water, quality of water in surrounding aquifers, well spacing, and needed pumpage during selected times). However, modeling the aquifer permits quantitative predictions, such as effect on ground-water levels due to limited surface-water irrigation in an area, effect on drain flows and river flow by ground-water pumpage, or optimum location of canals, wells, and drains in design of a system. It should be realized that modeling requires good basic-data input time to construct and verify the model, and realization of its limitations.

Studies by Conover (1954), Gunaji (1961), and Richardson (1971) along with streamflow measurements and chemical-quality analyses in the files of various agencies will provide some background information on ground water in the alluvium. The Bureau of Reclamation has records of monthly water-level measurements in 50 observation test holes located throughout the valleys. Richardson's digital model may be very useful in planning a basic-data collection program and even in later phases of the study as a management tool for operation of the irrigation system.

[p. 27] Components of the water-budget study are measured quantities of surface-water inflow, outflow, and precipitation; and calculated values of ground-water inflow, outflow, consumptive use by crops and phreatophytes, free-surface evaporation, and changes in storage.

In order to define quantities and quality of water a detailed program of basic-data collection and test-hole drilling will be necessary. The program would include such field work as well inventory, streamflow measurements of selected canals and drains, water-sampling of wells and surface flows for chemical analysis, and conducting aquifer tests and power-yield tests on selected wells. Some surface and borehole geophysical data may be collected.

A test-hole drilling program is needed to define the various quality-of-water zones laterally, and in depth. In addition to existing test holes and wells, as many as 25 additional shallow and cased test holes (± 80 foot depth) may be needed in order to sample the ground water in otherwise inaccessible locations.

At 15 or more sites throughout the Rincon and Mesilla Valleys it is recommended that "nests" of shallow, small-diameter sampling wells be installed in order to observe changes in the water-quality profile with time. These wells should be constructed as permanent observation and sampling test wells. Each "nest" would consist of 3 to 7 individually cased holes with depths ranging from about 10 to as much as 200 feet.

[p. 28] Two or three test holes ($\pm 1,500$ feet) should be drilled near the irrigated valleys in order to determine water quality and aquifer properties at depths below that which are normally encountered in wells. The exact location of test holes should be chosen after the well inventory is completed. These three deep test holes would be part of the recommended eight holes to be drilled and tested as part of the general study. Testing would include geophysical logging, water sampling, permeability tests, and hydrostatic-head measurements.

Using the basic data collected in the aforementioned programs, the hydrologic system can be analyzed and described in a preliminary written report. Based on the report a digital or combined analog-digital model would be constructed (or modified from an existing model). The model's purpose would be as a management tool, to show the hydrologic response to a given stress or imposed conditions on the irrigation system. It permits rapid calculation of quantitative effects of water-management practice[s].

The model would include river and flood-plain alluvium, as well as adjacent parts of the Santa Fe aquifer where heavy pumpage is affecting water in the alluvium (such as the Las Cruces well field). The model should be sensitive enough to duplicate changes in hydrologic conditions (such as water levels)

and small area of a few [sq. mi.] sections. Also, the model must be flexible enough to allow for large-scale modifications and changes in the present irrigation network.

[p. 29] Along with predicting changes in the quantities of water available at a particular location and time, the need is to predict the chemical quality of the water resulting from a management decision. It appears this second objective may also be solved by digital computer. The model may not have the sensitivity of the quantity model, but should show changes in ion concentration and loads over an area of several square miles during periods of a few years.

An open-file report would be written upon completion of the data gathering and defining of the system. Later, the modeling phase of the project, with supporting data, would be described in another report.

MAJOR ITEMS OF WORK [p. 30-33]

1. General collection of hydrologic data.
 - a. Well inventory - Inventory all wells in the upland area. Locate all large capacity wells in the flood-plain alluvium (estimated 2,000 wells) and inventory 40 percent or more on a selective basis. Inventory selected test holes: Take water samples of selected inventoried wells.
 - b. River, canal, and drain-flow measurements - The water-budget study will require frequent flow measurements and water sampling (weekly to monthly) of river, canal, and drain discharges at selected locations. The budget should follow the water year and continue over a 2- or 3-year period.
 - c. Aquifer tests - Pumping test and specific capacity measurements will be made whenever practical.
 - d. Pumpage inventory - Power-yield measurements need to be on wells in selected areas where acreage, irrigation quantities, and rainfall amounts are known. About 10 percent of the inventoried irrigation wells should be measured for the inventory.
 - e. Computer - formats will be used in data collection. Previously gathered material should be transferred to the format, and all format data put on punched cards and entered into a data file.
2. [p. 31] Quality-of-water work – Collect samples from 25 percent of all upland wells inventoried. Make specific-conductance measurements whenever possible. Sample, at selected depths, water from the deep test holes, and at intervals of time, water from the shallow test holes and observation wells. Later it may be desirable to install water-quality monitors on a few test holes.
3. Surface geophysics - a survey will be made of commercial sources, and if possible the data obtained. If desirable data are not available, a gravity survey of the area will be made by the Geological Survey. Minimum work would consist of five east-west gravity profiles distributed across the area, average length of 50 miles, using 1- mile station spacing. Altitudes of stations would be surveyed to an accuracy of 1 foot or less. In the vicinity of Las Cruces, more detailed gravity work is needed (estimated 75 additional stations). Based on analyses of the gravity survey, about 90 miles of seismic refraction profiles may be made at selected locations. Earth resistivity studies will be conducted in the Texas area of the Rio Grande Environmental Project area. If these studies prove successful, the seismic work may be replaced by resistivity studies.
4. [p. 32] Deep test drilling - Deep test drilling would be contracted to commercial drillers. Two or three holes are planned in the vicinity of Las Cruces, one ± 10 miles south of Las Cruces, two in the La Mesa bolson, two in the La Mesa bolson, one 10 to 20 miles west of Las Cruces; and one or two north of Las Cruces on the Jornada del Muerto. Location of these holes should depend on results of the geophysical surveys and well inventory. Each hole would be about 1,500 feet deep. Tests will be conducted at selected depth intervals in each hole to

determine (1) quality of water in that zone, (2) hydrostatic pressure, (3) aquifer permeability, and (4) lithology of the formation. If possible, in areas of high stress, each deep test hole will be cased and equipped so as to construct a permanent observation well where one or more separate intervals in the aquifers can be tested to monitor changes in water quality and hydrostatic pressure with time.

5. [p. 33] Shallow test-hole drilling - Drill about 25 shallow test holes in the alluvium as supplemental observation wells. At 15 other sites throughout the alluvium of the lower Rio Grande Valley, "nests" of observation wells would be installed. Depths of installation range from 10 to 200 feet, with depths of 10 to 80 feet being most common. These "nest" wells would be cased, perforated, and equipped as permanent observation wells. Two or three of the "nests" should be installed between the river alluvium and the city well field in order to monitor chemical changes with continuous city pumpage.
6. Borehole geophysical logging - borehole geophysical logs would be made in the eight deep test holes. All available logs on previously drilled wells and test holes would be purchased. The New Mexico District logger would make logs in shallow test holes, and in unused or abandoned deep test holes and wells. Also, some logging may be done by arrangement with commercial drillers.
7. Models - A digital or combined analog-digital model would be constructed for the area containing the flood-plain alluvium and Rio Grande. Also adjacent areas where heavy pumpage occurs would be included in the model. Exact planning on the models should await evaluation of the digital model used by Richardson (1971).

SCHEDULE OF WORK, TIME AND COSTS [p. 34-39]

First Fiscal Year, October 1971 – June

Schedule Item		Project Time	Costs
		In man-months	
1.	Project Planning		\$ 1,000
2.	Well Inventory		3,000
3.	Water-Use Data		2,000
4.	Water-Sample Collection		1,000
5.	Data Sources		4,000
6.	Water-Sample Analysis		2,000
7.	Report Preparation		4,000
8.	Report Review		1,200
9.	<u>Overhead Charge</u>		<u>1,800</u>
Totals		9	\$20,000

Second fiscal year, July 1972-June 1973

Schedule item	Project time in man-months	Costs
1. Begin inventory and sampling of wells. Put data on punch cards. (Estimated 2,800 wells in valley, inventory ±1,300 wells and sample 390 wells.)	13	\$ 53,690
2. Buy or do complete gravity geophysical survey.	1	8,200
3. Aerial magnetic geophysical survey.	-	6,000
4. Begin earth resistivity measurements or seismic profiling,	2	20,000
5. Plan and begin river, canal, and drain- flow measurements at selected sites. Begin water-budget calculations. Collect water samples for analyses.	5	17,500
6. Begin series of power-yield tests,	2	5,800
7. Drill, test, and equip shallow test holes in or near alluvium. About 25 single test holes (80' depth) and 15 "nests" of observation wells (10 to 200' depths). Work by USGS Hydrologic Laboratory, Denver, Sample wells bimonthly.	7	70,050
8. Anapra special report.	1	2,900
9. Aquifer tests on wells (pumping tests and specific capacity).	2	5,800
10. Prepare preliminary maps on geologic structure of base of Santa Fe Group and locate subsurface barriers.	1	2,900
11. Plan drilling of deep and shallow test holes,	1	2,900
12. Equipment to establish office.	-	3,000
13. Move allowance for 2 employees (or per diem).	-	3 000
14. Data analysis, progress reports, and project administration.	3	8,700
15. Drill 1 deep test hole (1,500').	<u>1</u>	<u>40,300</u>
TOTALS	39	\$250,740

Third fiscal year, July 1973-June 1974

Schedule item	Project time in man-months	Costs
1. Complete well inventory and sampling. Punch data.	4½	\$ 18,610
2. Complete earth resistivity and seismic work.	2	19,000
3. Measure and sample observation wells.	2½	21,880
4. Inventory new wells and update.	1	2,900
5. Make river, canal, and drain-flow measurements at selected sites, Collect and analyze water samples.	5	20,750
6. Continue aquifer tests.	3	8,700
7. Continue power-yield tests.	2	5,800
8. Drill 3 deep test holes (1,500') and test.	3	120,900
9. Begin detailed analysis of system, mainly in vicinity of Las Cruces. Prepare preliminary maps as possible, showing:		
a. Base of Santa Fe Group and subsurface structures and barriers.		
b. Major aquifers and/or water-quality zones.		
c. Altitude of water table.		
d. Base of alluvium and saturated thickness.		
e. Saturated sand thickness of different water-quality zones,		
f. Cross-sectional views of valley,	4	11,600
10. Assemble and review water budget.	3	7,900
11. Assemble and complete computer output of basic well data.	2	5,800
12. Data analysis, progress reports, and project administration.	4	11,600
TOTALS	36	\$255,440
36		

Fourth fiscal year, July 1974-June 1975

Schedule item	Project time in man-months	Costs
1. Drill 4 deep test holes (1,500'). Analyze drilling results.	6	\$167,000
2. Measure and sample observation wells.	2½	21,880
3. Inventory and update for new wells.	2	5,800
4. Make river, canal, and drain-flow measurements at selected sites. Collect and analyze water samples.	5	20,750
5. Continue aquifer tests.	2	5,800
6. Continue power-yield tests.	2	5,800
7. Borehole geophysical logging with District logger.	1	6,900
8. Outline contents of final reports. Plan digital models.	5	14,500
9. Continue intensive analysis of system. Refine water budget; complete maps started before; define geologic framework; begin water-quality budget.	4	11,600
10. Begin preparation of final reports and maps.	3½	10,150
11. Project administration, progress report, half-time secretary for ½ of year.	3 + 3 secretary	9,800
TOTALS	36 + 3 secretary	\$279,980

Fifth fiscal year, July 1975-June 1976		
Schedule item	Project time in man-months	Costs
1. Complete river, canal, and drain-flow measurements and sampling.	2	\$ 12,050
2. Measure and sample observation wells.	2	19,190
3. Complete power-yield tests and pumpage inventory.	1	2,900
4. Complete aquifer test program.	1	2,900
5. Inventory selected new wells and verify data.	2	8,260
6. Complete 2 open-file reports.	13	37,700
7. Complete digital model of ground water in alluvium.	4	11,600
8. Complete final report.	5	14,500
9. Secretary and project administration.	3 + 6 secretary	14,700
10. Effort on digital or analog-digital quality-of-water model.	3	8,700
TOTALS	36 + 6 secretary	\$132,500

[p. 39] A breakdown of expenses by year and cooperator is given below. It is assumed that the city of Las Cruces and Elephant Butte Irrigation District will share equal expenses throughout the last four years of the project. The Water Resources Research Institute furnishes office space to the project on a non-matching cooperative basis.

Costs					
State Engineer	City of Las Cruces	Elephant Butte Irrigation District	Anapra study	U.S. Geological Survey	Total
\$5,000	\$5,000			\$10,000	\$20,000
5,000	58,185	\$58,185	\$4,000	125,370	250,740
5,000	61,560	61,560	-	123,120	256,240
5,000	67,495	67,495		139,990	279,980
5,000	30,625	30,625		66,250	132,500
\$25,000	\$222,865	\$217,865	\$4,000	\$469,730	\$939,460

REPORTS [p. 40-41]

Progress reports would be made at the end of the second, third, and fourth fiscal years to all cooperators. Also at the end of the fourth fiscal year, a computer printout: of basic data such as a table of well records, results of chemical analyses, streamflow records and selected lithologic logs should be available. By January 1973, an open-file letter report with accompanying maps should be available on the general hydrologic conditions in the vicinity of the proposed port of entry at Anapra.

Two open-file reports, including maps and tables, would be prepared by about March 1976. One report would describe in detail the ground-water situation in the vicinity of Las Cruces. The other report would describe ground-water conditions in the flood-plain alluvium and adjacent aquifers of the Rio Grande Valley from Caballo Reservoir, New Mexico to El Paso, Tex. Some results of the quantitative model study should be included.

A comprehensive report to be completed by June 1976 would describe the general hydrologic situation of the southern Rio Grande Valley area, New Mexico. It is planned that this report would be published as a State Engineer Technical Report or as a Geological Survey Water-Supply Paper.

[p. 41] The final report from this project would contain results of the model studies on the flood-plain alluvium aquifer. This Report would describe the model and associated data used, verification, and the results obtained in using the model for design, planning, and operation of the surface-ground water system for irrigation in the Rincon and Mesilla Valleys. The date and type of report would depend on the results of the models. The completion date would be decided on later in the study.

SELECTED REFERENCES (p. 42-47)

- Basler, J.A., and Alary, L.J., 1968, Quality of shallow ground water in the Rincon and Mesilla Valley, New Mexico and Texas: U.S. Geological Survey Open-File Report 68-7, 30 p., 5 figs. **(G2, H2)**
- Blaney, J.A., and Hanson, E.G., 1965, Consumptive use and water requirements in New Mexico: New Mexico State Engineer Technical Report 32, 32 p., 6 figs. **(D1)**

- Conover, C.S., 1954, Ground-water conditions in the Rincon and Mesilla Valleys and adjacent areas in New Mexico: U.S. Geological Survey Water-Supply Paper 1230, 200 p. **(G2)**
- Conover, C.S., Herrick, E.H., Wood, J.W., and Weir, J.E., Jr., 1955, The occurrence of ground water in south-central New Mexico: New Mexico Geological Society Guidebook 5, p. 108-119. **(G2)**
- Córdoba, D.A., Wengerd, S.A., and Shomaker, J.W., eds., 1969, Guidebook of the Border Region: New Mexico Geological Society Guidebook 20, 218 p. *See Road Log Section, p. 1-38.* **(C2a)**
- Cox, E.R., and Reeder, H.O., 1962, Ground-water conditions in the Rio Grande Valley between Las Palomas and Truth or Consequences, New Mexico: N.M. State Engineer Technical Report 25, 47 p. **(G2, H1)**
- Davis, L.V., and Busch, F.E., 1968, Summary of hydrologic investigations by the United States Geological Survey at White Sands Missile Range, New Mexico: U.S. Geological Survey Open-File Report 68-70, 299 p., 27 figs. **(G2)**
- Davis, M.E., 1962, Development of ground water in the El Paso District, Texas 1960-63, Progress Report No. 9: Texas Water Commission Bulletin 6514, 34 p., 9 figs. **(G2)**
- Dinwiddie, G.A., Mourant, W.A., and Basler, J.A., 1966, Municipal water supplies and uses, southwestern New Mexico; New Mexico State Engineer Technical report 29D, Santa Fe, New Mexico, p. 17-34. **(E2)**
- Dixon, G.H., Baltz, E.H., Stipp, T.F., and Bieberman, R.A., 1955, Map of New Mexico showing test wells for oil and gas, oil and gas fields, and pipelines: U.S. Geological Survey Oil and Gas Investigations Map OM-159. **(C2a)**
- Doty, G.C., 1963, Water-supply development at the National Aeronautics and Space Agency-Apollo Propulsion System Development Facility, Doña Ana County, New Mexico: U.S. Geological Survey Open-File Report 63-29, 40 p. **(C2a, H1)**
- Dunham, K.C., 1935, The geology of the Organ Mountains, with an account of the geology and mineral resources of Dona Ana County, New Mexico: New Mexico Bureau of Mines and Mineral Resources, Bulletin 11, 272 p. **(C2a, G1)**
- Feth, J.H., 1965a, Preliminary map of the conterminous United States showing depth to and quality of shallowest ground water containing more than 1,000 parts per million dissolved solids: U.S. Geological Survey Hydrologic Atlas 199, 31 p., 2 pls., 1:3,000,000. **(D1, H2)**
- Hawley, J.W., 1965, Geomorphic surfaces along the Rio Grande Valley from El Paso, Texas to Caballo Reservoir, New Mexico: New Mexico Geological Society, Guidebook 16, p. 188-198. **(C2a)**
- Hawley, J.W., compiler, 1970, Cenozoic stratigraphy of the Rio Grande Valley area, Doña Ana County, New Mexico: El Paso Geological Society, Guidebook 4, 49 p. **(C2a, F2)**
- Hawley, J.W., and Gile, L.H., 1966, Landscape evolution and soil genesis in the Rio Grande region, southern New Mexico: Friends of the Pleistocene, Rocky Mountain Section, Guidebook 11th Field Conference; La Cruces, New Mexico State University Agronomy Department, Special Publication, 74 p. **(C2a, C3)**
- Hawley, J.W., and Kottlowski, F.E., 1969, Quaternary geology of the south-central New Mexico border region, *in* Border Stratigraphy Symposium: New Mexico Bureau of Mines and Mineral Resources, Circular 104, p. 89-115. **(C2a)**
- Hawley, J.W., Kottlowski, F.E., Seager, W.R., King, W.E., Strain, W.S., and LeMone, D.V., 1969, The Santa Fe Group in the south-central New Mexico border region, *in* Border Stratigraphy Symposium: New Mexico Bureau of Mines and Mineral Resources, Circular 104, p. 52-76. **(C2a, F1)**
- Herrick, E.H., and Davis, L.V., 1965, Availability of ground water in the Tularosa Basin and adjoining areas: U.S. Geological Survey Hydrologic Investigations Atlas, HA-191, scale 1:760,320. **(D1, G2)**
- Keyes, C.R., 1905, Geology and underground water conditions of the Jornada del Muerto, New Mexico: U.S. Geological Survey Water-Supply and Irrigation Paper 123, 42 p. **(G1)**
- King, W.E., Hawley, J.W., Taylor, A.M., and Wilson, R.P., 1969, Hydrogeology of the Rio Grande Valley and adjacent intermontane areas of southern New Mexico: New Mexico Water Resources Research Institute Miscellaneous Report No. M6, 141 p. <https://nmwrri.nmsu.edu/publications/miscellaneous-reports/m-documents/m6.pdf> **(F2, H1, H2)**
- King, W.E., Hawley, J.W., Taylor, A.M., and Wilson, R.P., 1971, Geology and ground-water resources of central and western Doña Ana County, New Mexico: New Mexico Bureau of Mines and Mineral Resources, Hydrologic Report 1, 64 p. **(F2, H1, H2)**
- Kirby, J.W., 1968, Water resources – El Paso County, Texas, past, present, future: University of Texas at El Paso, master's thesis, 190 p., 5 figs. **(F2, G2)**
- Kottlowski, F.E., 1955, Cenozoic sedimentary rocks in south-central New Mexico: New Mexico Geological Society, Guidebook 6, p. 88-91. **(C2a)**

- Kottlowski, F.E., 1965a, Facets of the Late Paleozoic strata of southwestern New Mexico: New Mexico Geological Society Guidebook 16, p., 141-147. **(C2a)**
- Kottlowski, F.E., 1965b Sedimentary basins of south-central and southwestern New Mexico: American Association of Petroleum Geologists Bulletin, v. 49, nos. 11 and 12, p. 2120-2139. **(C2a)**
- Kottlowski, F.E., Flower, R.H., Thompson, M.L., and Foster, R.W., 1956, Stratigraphic studies of the San Andres Mountains, New Mexico: New Mexico Bureau of Mines and Mineral Resources, Memoir 1, 132 p. **(C2a)**
- Lee, W.T., 1907, Water resources of the Rio Grande Valley in New Mexico: U.S. Geological Survey Water-Supply Paper 188, 59 p. **(C2a, F2, G1)**
- Leggat, E.R., 1962, Development of ground-water of the El Paso district, Texas: 1955-60, Progress Report No. 8: Texas Water Commission, Bulletin 6204, 56 p. **(F2, G2)**
- Leggat, E.R., Lowry, M.E., and Hood, J.W., 1962, Ground-water resources of the lower Mesilla Valley, Texas and New Mexico: Texas Water Commission, Bulletin 6203, 191 p. **(F2, G2, H1, H2)**
- Maker, H.J., Neher, R.E., Derr, P.H., and Anderson, J.U., 1971, Soil Associations and land classifications for irrigation, Doña Ana County: New Mexico State University Agricultural Experiment Station, Research Report 183, 40 p. **(C3)**
- Meyer, W.R., and J.D. Gordon, 1972, Development of ground water in the El Paso district, Texas, 1963–1970, *in* Texas Water Development Board Report 53, 50 p. **(H3)**
- Nelson, L.A., 1940, Paleozoic stratigraphy of the Franklin Mountains of West Texas: American Association of Petroleum Geologists Bulletin, v. 24, no. 1, p. 157-172. **(C2a)**
- Richardson, G.L., 1971, Water-table investigation in the Mesilla Valley, unpublished Master's thesis, New Mexico State University. *See Richardson and others, 1972.* **(H3)**
- Richardson, G.L., Gebbard, T.G., Jr., and Brutsaert, W.F., 1972, Water-table investigation in the Mesilla Valley: Las Cruces, New Mexico State University, Engineering Experiment Station Technical Report 76, 206 p. *See Richardson 1971.* **(H3)**
- Ruhe, R.V., 1967, Geomorphic surfaces and surficial deposits in southern New Mexico: New Mexico Bureau of Mines and Mineral Resources, Memoir 18, 65 p. **(C2a, C3)**
- Slichter, C.S., 1905, Observations on ground waters of the Rio Grande valley: U.S. Geological Survey Water-Supply and Irrigation Paper 141, 83 p. **(G1)**
- Smith, R.E., 1956, Ground water of the El Paso District, Texas: Texas Board of Water Engineers, Bulletin 5603, 33 p. **(C2a, F1, G2)**
- U.S. Geological Survey, 1968a, Water resources data for New Mexico, part 1. Surface-water records: U.S. Geological Survey Water Resources Division, Albuquerque, 242 p., 4 figs. **(D1)**
- U.S. Geological Survey, 1968b, Water resources data for New Mexico, part 2. water quality records: U.S. Geological Survey Water Resources Division, Albuquerque, 207 p., 2 figs. **(D1, E2a)**
- Vlissides, S.D., and Bieberman, R.A., 1961, Map of New Mexico showing oil and gas fields, unsuccessful test wells, Precambrian rocks, and pipelines: U.S. Geological Survey Oil and Gas Investigations Map OM-207. **(C2a)**